

(19) World Intellectual Property Organization
International Bureau



PCT



(43) International Publication Date
26 May 2006 (26.05.2006)

(10) International Publication Number
WO 2006/054177 A1

(51) **International Patent Classification:**
C12N 15/11 (2006.01) **A61P 35/00** (2006.01)

(21) **International Application Number:**
PCT/IB2005/003641

(22) **International Filing Date:**
18 November 2005 (18.11.2005)

(25) **Filing Language:** English

(26) **Publication Language:** English

(30) **Priority Data:**
PCT/IB2004/004093
19 November 2004 (19.11.2004) IB
11/144,322 3 June 2005 (03.06.2005) US

(71) **Applicant** (for all designated States except US): **INSTITUT GUSTAVE ROUSSY** [FR/FR]; 39, rue Camille Desmoulins, F-94805 Villejuif Cedex (FR).

(72) **Inventors; and**

(75) **Inventors/Applicants** (for US only): **ANDRE, Fabrice** [FR/FR]; 6, rue Veronese, F-75013 Paris (FR). **ZITVOGEL, Laurence** [FR/FR]; 17, rue Nord, F-92160 Antony (FR). **SABOURTN, Jean-Christophe** [FR/FR]; 298, Rue Charenton, F-75012 Paris (FR).

(74) **Agents:** **GALLOIS, Valérie** et al.; Becker et Associates, 25, rue Louis le Grand, F-75002 Paris (FR).

(81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW

(84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, **DK**, EE, ES, FI, FR, GB, GR, HU, IE, **IS**, IT, LT, LU, LV, MC, NL, PL, PT, **RO**, SE, **SI**, SK, TR), OAPI (BF, **BJ**, CF, CG, CI, CM, GA, GN, GQ, GW, ML, **MR**, NE, SN, TD, TG).

Declaration under Rule 4.17:

— of inventorship (Rule 4.17(iv))

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) **Title:** IMPROVED TREATMENT OF CANCER USING TLR3 AGONISTS

(57) **Abstract:** The present invention relates generally to the fields of genetics and medicine. More specifically, the present invention relates to improved methods of treating cancers using a TLR3 agonist, by assessing the expression of a TLR3 receptor by cancer cells.



WO 2006/054177 A1

IMPROVED TREATMENT OF CANCER USING TLR3 AGONISTS

TECHNICAL FIELD OF THE INVENTION

5 The present invention relates generally to the fields of genetics and medicine. More specifically, the present invention relates to improved methods of treating cancers using TLR3 agonists.

BACKGROUND OF THE INVENTION

10 Cancer is one of the leading causes of death in Europe and in the United States. There are more than 1.2 million cancer cases each year in the U.S. and cancer is expected to be the leading cause of the death by the year 2010. Lung and prostate cancer are the top cancer killers for men in the United States. Lung and breast cancer are the top cancer killers for women in the United States. A cure for cancer has yet to be found. Current treatment options, such as surgery, chemotherapy and
15 radiation treatment, are oftentimes either ineffective or present serious side effects.

Currently, cancer therapy may involve surgery, chemotherapy, hormonal therapy and/or radiation treatment to eradicate neoplastic cells in a patient. Recently, cancer therapy could also involve biological therapy or immunotherapy. All of these approaches pose significant drawbacks for the patient. Surgery, for example, may be contraindicated due to the health of the patient or may be
20 unacceptable to the patient. Additionally, surgery may not completely remove the neoplastic tissue.

Despite the availability of a variety of chemotherapeutic agents, chemotherapy has many drawbacks. Almost all chemotherapeutic agents are toxic, and chemotherapy causes significant, and often dangerous, side effects, including severe nausea, bone marrow toxicity,
25 inummosuppression, etc. Additionally, even with administration of combinations of chemotherapeutic agents, many tumor cells are resistant or develop resistance to the chemotherapeutic agents. In fact, those cells resistant to the particular chemotherapeutic agents used in the treatment protocol often prove to be resistant to other drugs, even those agents that act by mechanisms different from the mechanisms of action of the drugs used in the specific treatment;
30 this phenomenon is termed pleiotropic drug or multidrug resistance. Thus, because of drug resistance, many cancers prove refractory to standard chemotherapeutic treatment protocols.

Double-stranded RNA molecules, such as poly A-polyU and poly I-poly U, are immunostimulating agents. Preclinical studies performed in 1970-1980's showed that the incubation of blood mononuclear cells with poly A-poly U induces interferon alpha secretion, and that the injection of
35 poly A-poly U activates natural killer cells in vitro (EP281 380 ; EP 113 162).

It has recently been demonstrated that the double-stranded RNA receptor is Toll Like receptor 3 (TLR3) (Alexopoulou et al, 2001). This receptor has been described to be expressed in membranes

of dendritic cells and of cells from colic mucosa. The binding of double-stranded RNA to this receptor activates dendritic cells and activates T lymphocytes. Consequently, the use of double-stranded RNA for treating cancer has been attempted. However, the response rate was not high and no therapeutic applications were developed.

- 5 Therefore, a method allowing to select responding patient would greatly enhance the therapeutic efficacy of double-stranded therapies.

SUMMARY OF THE INVENTION

- 10 The present invention demonstrates the existence of a correlation between the expression of a TLR3 in cells from a tumor sample in a subject and the ability of said subject to respond to treatment with a composition comprising a TLR3 agonist, preferably an antibody or other TLR3 binding protein, a phosphate containing small molecule, an organophospho-ester, a nucleotide, a nucleotide-like compound, a nucleotide analog, or a nucleic acid molecule, or most preferably a
- 15 double-stranded RNA. More specifically, the present invention shows, for the first time, that TLR3 is expressed in tumoral cell membranes and that the binding of a TLR3 agonist, in particular double-stranded RNA, on said tumoral cells through the TLR leads to tumoral cells lysis and tumor regression. In contrast, tumoral and other cells that do not express TLR are not sensitive to a TLR3 agonist treatment, in particular the double-stranded RNA treatment. The finding is particularly
- 20 surprising and important for the management of cancer patients because an increase in survival is rarely if ever achieved in any known active regimen in metastatic breast cancer. The usual finding is an improvement in early endpoints (response rate and/or time to progression, otherwise known as progression-free survival) but the improvement almost never translates into a significant increase in overall survival, as exhibited in patients treated in accordance with the present invention.
- 25 Furthermore, the finding that TLR3 agonist therapy can increase survival will permit therapeutic strategies that are expected to be efficacious and a suitable treatment for tumors, including particularly breast tumors, whether they involve axillary lymph nodes or not, that is beyond only breast cancers presenting axillary lymph node involvement. Likewise, the TLR3 agonists may find use in breast (and other) cancers that are metastatic, recurring and/or refractory.
- 30 Accordingly, in another aspect, the present invention provides a method of treating a human subject having a tumor comprising TLR3-expressing cells, which method comprises administering to said patient a therapeutically effective amount of at least one TLR3 agonist. In a preferred embodiment, said human patient has a solid tumor. In a preferred embodiment, said human patient has a breast cancer.
- 35 Therefore, the present invention concerns the use of a TLR3 agonist for the manufacture of a medicament for treating cancer in a subject, wherein said cancer in said subject comprises cancer cells expressing a TLR3 receptor. More preferably, the TLR3 agonist is a double-stranded RNA molecule. Optionally, the cancer is a metastatic cancer. In a particular embodiment, the present

invention concerns the use of a double-stranded polyA/polyU RNA molecule for the manufacture of a medicament for treating a cancer in a subject, wherein said cancer in said subject comprises cancer cells expressing a TLR3 receptor. Preferably the cancer is a solid tumor or a carcinoma, for example a breast cancer. The present invention also concerns a method for assessing the response of
5 a subject having cancer to a treatment using a TLR3 agonist, the method comprising determining whether cancer cells in said subject express a TLR3 receptor, the expression of a TLR3 receptor being indicative of a responder subject. More preferably, the TLR3 agonist is a double-stranded RNA molecule.

The present invention further concerns a method for selecting subjects having a cancer that respond
10 to a treatment using a TLR3 agonist, the method comprising determining whether cancer cells in said subject express a TLR3 receptor, the expression of a TLR3 receptor being indicative of a responder subject. More preferably, the TLR3 agonist is a double-stranded RNA molecule.

In addition, the present invention concerns a method for treating a subject having a cancer, the method comprising determining whether cancer cells in said subject express a TLR3 receptor, the
15 expression of a TLR3 receptor being indicative of a subject responding to a TLR3 agonist, and treating said subject whose cancer cells express a TLR3 receptor with a TLR3 agonist. More preferably, the TLR3 agonist is a double-stranded RNA molecule. In a related embodiment, the method for treating a subject having a cancer comprises the additional step of surgically removing a portion of the cancer prior to treating said subject with a TLR3 agonist. In yet another
20 embodiment, the method comprises the additional step of treating said patient with radiotherapy in combination with a TLR3 agonist.

In another embodiment, the invention provides a method for treating a subject having a cancer, comprising the steps of: administering to said subject an agent that causes increased expression of a TLR3 receptor in cancer cells; and administering to said patient a TLR3 agonist. In a related
25 embodiment, this method of treatment additionally comprises the step of determining whether cancer cells in said subject express a TLR3 receptor. This determination step may be performed subsequent to and/or following the treatment with the agent that causes increased expression of TLR3 receptor, but prior to treating said subject with a TLR3 agonist. In another related embodiment, this method comprises the additional step of surgically removing a portion of the
30 cancer prior to administering the agent that causes increased expression of a TLR3 receptor. In yet another embodiment, the method comprises the additional step of treating said patient with radiotherapy in combination with a TLR3 agonist.

Moreover, the inventors have surprisingly found in the follow up study to the 300 patient clinical trial that patients having a solid tumor and treated with a TLR3 agonist in accordance with a repeat-
35 dose and with an intravenous therapeutic regimen exhibited greater survival than other patients not treated with the TLR3 agonist.

Accordingly, in one aspect, the present invention provides a method of achieving enhanced or prolonged survival in human patients with cancer, which comprises administering to said patient a therapeutically effective amount of at least one TLR3 agonist. In a preferred embodiment, said at least one TLR3 agonist is intravenously administered. In a preferred embodiment, the cancer is a breast cancer. In a more particular embodiment, the cancer is a metastatic or recurrent breast cancer. In another preferred embodiment, the cancer is a node positive breast cancer. La yet another preferred embodiment the breast cancer involved 1 to 3 lymph nodes.

Accordingly, in one aspect, the present invention provides a method of achieving enhanced survival in human patients with cancer or a method for treating a subject having a cancer, which comprises administering to said patient (a) at least a first dose therapeutically effective amount of at least one TLR3 agonist; and (b) at least a second dose of a therapeutically effective amount of at least one TLR3 agonist. In a preferred embodiment, said first and said second doses are administered at an interval of less than one month, less than three weeks, less than two weeks, or less than one week. In another preferred embodiment, the subject is administered an intravenous dose of a TLR3 agonist about once a week for a period of six weeks. In another preferred embodiment, the TLR3 agonist is a double-stranded RNA molecule. In another preferred embodiment, the TLR3 agonist is selected from polyA:polyU or polyLpolyC. In a more preferred embodiment, the dose of polyA:polyU or polyLpolyC is between about 10mg and 100 mg.

In a preferred embodiment of the methods and uses according to the present invention, the subject is a human subject.

It will be appreciated that the methods of treatment mentioned herein can be used as prophylactic treatment; in any of the embodiments herein, a prophylactically effective amount of the TLR3 agonist can be interchanged with a therapeutically effective amount of a TLR3 agonist.

In a preferred embodiment of the methods and uses according to the present invention, the cancer is a solid tumor or a carcinoma. Preferably, the solid tumor is selected from breast cancer, colon cancer, lung cancer, prostate cancer, renal cancer, metastatic or invasive malignant melanoma, brain tumor, bladder cancer and liver cancer. Carcinoma includes bladder, breast, colon, kidney, liver, lung, ovary, pancreas, stomach, cervix, thyroid or skin carcinoma, including squamous cell carcinoma. In a most preferred embodiment, the solid tumor is a breast cancer. However, the present invention also contemplates hematopoietic tumors such as leukemia, acute lymphocytic leukemia, acute lymphoblastic leukemia, B-cell lymphoma, T-cell lymphoma, Hodgkin's lymphoma, non-Hodgkin's lymphoma, hairy cell lymphoma, Burkett's lymphoma, acute and chronic myelogenous leukemias and promyelocyte leukemia. The present invention is also relevant for the treatment of metastasis.

In a preferred embodiment, the expression of a TLR3 receptor in said cancer cell is determined using a TLR3-specific ligand. Preferably, the ligand is an antibody, or a fragment or derivative thereof.

In an alternative embodiment, the expression of a TLR3 receptor in said cancer cell is determined using a TLR3-specific primer or probe.

Preferably, the expression of a TLR3 receptor in said cancer cell is determined in vitro or ex vivo. However, the determination in vivo is also encompassed by the present invention.

5 In a preferred embodiment of the methods and uses according to the present invention, the double-stranded RNA molecule is a polyA/polyU ("polyAU") molecule. In an other preferred embodiment of the methods and uses according to the present invention, the double-stranded RNA molecule is a polyI/polyC molecule.

The present invention further concerns a kit for selecting subjects that respond to a treatment using
10 a TLR3 agonist, more preferably a double-stranded RNA molecule, the kit comprising reagents for determining the expression of a TLR3 receptor in a cancer cell in a sample. In yet another aspect, the invention relates to compositions of matter and therapeutic and screening methods that take advantage of the presence of TLR3 on a cancer cell surface. In one embodiment, the invention provides a complex comprising an agent that binds to TLR3 in association with a tumoricidal or
15 cytotoxic agent. In another embodiment, the invention provides a method of treating a subject with a cancer characterized by the presence of TLR3 on a cancer cell surface comprising the step of administering to said patient such a complex. In another embodiment, the invention provides a method of determining if a test compound is useful for the treatment of cancer, said method comprising the step of determining if said test compound is a TLR3 agonist. In a preferred
20 embodiment, the method of determining if a test compound is useful for the treatment of cancer comprises the steps of: contacting said test compound with a cancer cell characterized by the presence of TLR3 on its cell surface; and determining if said test compound induces a TLR3-mediated biological effect. Such a method is useful for screening proteins, small molecules, nucleic acids and other biomolecules.

25

DESCRIPTION OF THE FIGURES

Figure 1 illustrates the TLR3 expression by primary tumor. TLR3 is overexpressed by tumor cells in 10% of samples (n=18).

Figure 2 illustrates Survival of patients with TLR3- tumors (figure 2a) or with TLR3+ tumors
30 (figure 2b) according to treatment with a placebo (observation) or with dsRNA.

DETAILED DESCRIPTION OF THE INVENTION

A marker-based approach to tumor identification and characterization promises improved diagnostic and prognostic reliability. Typically, the diagnosis of breast cancer and other types of
35 cancer requires histopathological proof of the presence of the tumor. In addition to diagnosis, histopathological examinations also provide information about prognosis and selection of treatment regimens. Prognosis may also be established based upon clinical parameters such as tumor size,

tumor grade, the age of the patient, and lymph node metastasis.

With the available and potent conventional drug regimens as well as the advent of novel therapy approaches targeting specific biological pathways, the determination of optimal treatment of a primary cancer is becoming increasingly complex. Moreover, the outcome of a treatment of a

5 patient with cancer is often unpredictable. Only a portion of the patients respond to a certain type of treatment. The patients receiving a specific type of treatment are subjected to an unnecessary suffering since adverse reactions often are obtained from certain treatment used. Some treatments elicit more severe reaction from the patient than other treatments. Mostly, the effect of a treatment is not shown until 3-6 months after treatment. It would therefore be of great importance if patients
10 with a high probability to respond could be identified before the onset of treatment. To date, no set of satisfactory predictors for prognosis or therapeutic response based on the clinical information alone has been identified for TLR agonist compounds in cancer therapy.

As further described herein, the studies disclosed herein present results from 300 patients with early breast cancer that had been included from 1972 to 1979 in a randomized trial comparing post-
15 operative administration of TLR agonist polyAU with no treatment. When the TLR was investigated as a potential biomarker, it was observed that tumor biopsies positive for TLR demonstrated high long term survival rates, and presumably that this survival is in response to therapy with the TLR3 agonist. Patient biopsies were stained with a TLR3-specific monoclonal antibody ("mAb") and correlation of TLR3 expression with polyAU efficacy was determined. 182
20 tumor samples (91%) were available from the 200 patients included in the randomized trial. TLR3 was strongly expressed by tumor cells in 18 patients (10%). TLR3 expression correlated with a significantly increased 20-year survival rate.

Accordingly, in one aspect, the present invention provides a method of determining whether a patient will respond to therapy with a TLR3 agonist. In one embodiment the present invention
25 concerns a method for selecting or identifying a subject having a tumor that will respond to a treatment using a TLR3 agonist, the method comprising determining whether tumor cells in said subject express a TLR3, the expression of a TLR3 receptor being indicative of a responder subject. In a preferred embodiment, expression of TLR3 is determined using an immunohistochemical assay, as used in the Examples. Preferably TLR3 expression is determined using an antibody that
30 specifically binds TLR3.

In another embodiment, the present invention provides a method for characterizing a cell or a tumor in a patient, the method comprising: obtaining or providing a biological sample from the patient, wherein said sample comprises a tumor cell or biological material derived therefrom, and determining whether said cell expresses a TLR3. In a preferred embodiment, the present invention
35 provides a method for characterizing a cell or a tumor in a patient, the method comprising: obtaining or providing a tumor biopsy from the patient, and determining whether said biopsy

comprises a cell expressing a TLR3 polypeptide.

DEFINITIONS

As used in the specification, "a" or "an" may mean one or more. As used in the claim(s), when used
5 in conjunction with the word "comprising", the words "a" or "an" may mean one or more than one.
As used herein "another" may mean at least a second or more.

Where "comprising" is used, this can preferably be replaced by "consisting essentially of," more preferably by "consisting of."

The terms "cancer" and "tumor" as used herein are defined as a new growth of cells or tissue
10 comprising uncontrolled and progressive multiplication. In a specific embodiment, upon a natural course the cancer is fatal. In specific embodiments, a cancer is invasive, metastatic, and/or anaplastic (loss of differentiation and of orientation to one another and to their axial framework).

The term "breast cancer" as used herein is defined as cancer which originates in the breast. In a specific embodiment, the breast cancer has spread to other organs, such as lymph nodes. In a
15 specific embodiment, the breast cancer is invasive and may be metastatic.

The term "invasive" as used herein refers to cells which have the ability to infiltrate surrounding tissue. In a specific embodiment, the infiltration results in destruction of the surrounding tissue. In another specific embodiment, the cells are cancer cells. In a preferred embodiment, the cells are breast cancer cells, and the cancer has spread out of a duct into surrounding breast epithelium. In a
20 specific embodiment, "metastatic" breast cancer is within the scope of "invasive."

The term "metastatic" as used herein is defined as the transfer of cancer cells from one organ or part to another not directly connected with it. In a specific embodiment, breast cancer cells spread to another organ or body part, such as lymph nodes.

"Weekly" stands for "about once a week" (meaning that more than one treatment is made with an
25 interval of about one week between treatments), the about here preferably meaning +/-1 day (that is, translating into "every 6 to 8 days"); most preferably, "weekly" stands for "once every 7 days".

The term "biopsy" as used herein is defined as removal of a tissue from an organ (e.g., breast) for the purpose of examination, such as to establish diagnosis. Examples of types of biopsies include by application of suction, such as through a needle attached to a syringe; by instrumental removal
30 of a fragment of tissue; by removal with appropriate instruments through an endoscope; by surgical excision, such as of the whole lesion; and the like.

As used herein, the term "adjunctive" is used interchangeably with "in combination" or "combinatorial". Such terms are also used where two or more therapeutic or prophylactic agents affect the treatment or prevention of the same disease. For the avoidance of doubt, two agents used
35 "in combination" may be, but are not necessarily, co-administered.

As used herein, the term "in combination" refers to the use of more than one therapies (e.g., more than one prophylactic agent and/or therapeutic agent). The use of the term "in combination" does

not restrict the order in which therapies (e.g., prophylactic or therapeutic agents) are administered to a subject with cancer. A first therapy can be administered prior to (e. g. , 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12 hours, 24 hours, 48 hours, 72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks before),
5 concomitantly with, or subsequent to (e. g., 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12 hours, 24 hours, 48 hours, 72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks after) the administration of a second therapy to a subject with cancer.

By "co-administration" or "co-administering" we mean that the two agents are administered in
10 temporal juxtaposition. The co-administration may be effected by the two agents being mixed into a single formulation, or by the two agents being administered separately but simultaneously, or separately and within a short time of each other. For example, in general the two agents are co-administered within the time range of 12-72 hours. In this case, the agents may be administered in either order. In a preferred embodiment of the instant invention, the two agents are co-administered
15 in a single formulation, or are co-administered simultaneously.

As used herein, the terms "prevent", "preventing" and "prevention" refer to the inhibition of the development or onset of cancer (particularly, a breast cancer) or the prevention, recurrence, onset, or development of one or more symptoms of cancer, particularly a breast cancer, in a subject resulting from the administration of therapy (e.g., a prophylactic or therapeutic agent) or a
20 combination of therapies (e.g., a combination of prophylactic and/or therapeutic agents).

As used herein, the term "prophylactically effective amount" refers to that amount of the prophylactic agent sufficient to result in the prevention of the recurrence or onset of cancer or one or more symptoms thereof.

As used herein, a "protocol" includes dosing schedules and dosing regimens. The protocols herein
25 are methods of use and include prophylactic and therapeutic protocols.

As used herein, a "prophylactic protocol" refers to a regimen for dosing and timing the administration of one or more prophylactic agents.

As used herein, the term "therapeutic protocol" refers to a regimen for dosing and timing the administration of one or more therapeutic agents.

30 As used herein, the phrase "side effects" encompasses unwanted and adverse effects of a therapy (e.g., prophylactic and/or therapeutic agent). Adverse effects are always unwanted, but unwanted effects are not necessarily adverse. An adverse effect from a prophylactic or therapeutic agent might be harmful or uncomfortable or risky.

As used herein, the term "small molecules" includes, but is not limited to, organic or inorganic
35 compounds (i.e., including heteroorganic and organometallic compounds) having a molecular weight less than 1,000 grams per mole, hi a preferred embodiment, "small molecules" encompass organic or inorganic compounds having a molecular weight less than 750 grams per mole. In yet

another specific embodiment, "small molecules" encompass organic or inorganic compounds having a molecular weight less than 500 grams per mole. Salts, esters, and other pharmaceutically acceptable forms of such compounds are also encompassed.

As used herein, the terms "subject" and "patient" are used interchangeably.

5 As used herein, the terms "subject" and "subjects" refer to an animal, preferably a mammal including, but not limited to, a non-primate (e.g., a cow, pig, horse, cat, dog, rat, and mouse) and a primate (e.g., a monkey such as a cynomolgous monkey and a human), and more preferably a human. In a specific embodiment, the subject is a human with cancer. In a preferred embodiment, the subject is a human with a solid tumor, for example a breast cancer.

10 As used herein, the term "synergistic" refers to a combination of therapies (e.g., prophylactic or therapeutic agents) which is more effective than the additive effects of any two or more single agents. For example, a synergistic effect of a combination of therapies (e. g. , prophylactic or therapeutic agents) permits the use of lower dosages of one or more of the agents and/or less frequent administration of said therapies to a subject with cancer.

15 The ability to utilize lower dosages of therapies (e. g. , prophylactic or therapeutic agents) and/or to administer said therapies less frequently reduces the toxicity associated with the administration of said therapies to a subject without reducing the efficacy of said therapies in the prevention or treatment of cancer. In addition, a synergistic effect can result in improved efficacy of therapies in the prevention or treatment of cancer. Finally, synergistic effect of a combination of therapies may
20 avoid or reduce adverse or unwanted side effects associated with the use of any single therapy.

As used herein, the terms "therapeutic agent" and "therapeutic agents" refer to any agent (s) which can be used in the treatment, management, or amelioration of cancer or one or more symptoms thereof. Preferably, a therapeutic agent is an agent which is known to be useful for, or has been or is currently being used for the treatment, management, or amelioration of cancer or one or more
25 symptoms thereof. A single therapeutic agent may be characterized as two or more different types of agents based upon one or more effects the agent has in vivo and/or in vitro.

As used herein, the term "effective amount" refers to the amount of a therapy (e. g. a prophylactic or therapeutic agent) which is sufficient to reduce or ameliorate the severity, duration and/or progression of cancer or one or more symptoms thereof, ameliorate one or more symptoms of
30 cancer, prevent the advancement of cancer, cause regression of cancer, prevent the recurrence, development, or onset of cancer or one or more symptoms thereof, or enhance or improve the prophylactic or therapeutic effect (s) of another therapy (e. g., prophylactic or therapeutic agent).

As used herein, the term "therapeutically effective amount" refers to that amount of a therapy (e. g., a therapeutic agent) which is sufficient to destroy, modify, control or remove primary, regional or
35 metastatic cancer tissue, ameliorate cancer or one or more symptoms thereof, or prevent the advancement of cancer, cause regression of cancer, or enhance or improve the therapeutic effect (s) of another therapy (e. g., a therapeutic agent).

- A therapeutically effective amount may refer to the amount of a therapy (e.g. , a therapeutic agent) sufficient to delay or minimize the spread of cancer. A therapeutically effective amount may also refer to the amount of a therapy (e. g. , a therapeutic agent) that provides a therapeutic benefit in the treatment or management of cancer. Further, a therapeutically effective amount with respect to a
- 5 therapeutic agent of the invention means that amount of therapeutic agent alone, or in combination with other therapies, that provides a therapeutic benefit in the treatment or management of cancer. Used in connection with an amount of a TLR3 agonist, the term can encompass an amount that improves overall therapy, reduces or avoids unwanted effects, or enhances the therapeutic efficacy of or synergizes with another therapy (e. g. , a therapeutic agent).
- 10 As used herein, the terms "treat", "treatment" and "treating" refer to the reduction or amelioration of the progression, severity, and/or duration of cancer, particularly a solid tumor, for example a breast cancer, or one or more symptoms thereof that results from the administration of one or more therapies (e.g., one or more prophylactic and/or therapeutic agents).
- As used herein, the terms "therapies" and "therapy" can refer to any protocol(s), method(s) and/or
- 15 agent(s) that can be used in the prevention, treatment, management or amelioration of cancer or one or more symptoms thereof. In certain embodiments, the terms "therapy" and "therapies" refer to cancer chemotherapy, radiation therapy, hormonal therapy, biological therapy, and/or other therapies useful for the prevention, management, or treatment of cancer known to an oncologist skilled in the art.
- 20 As used herein, the terms "manage", "managing" , and "management" refer to the beneficial effects that a subject derives from a therapy (e. g. , a prophylactic or therapeutic agent), which does not result in a cure of the disease, hi certain embodiments, a subject is administered one or more therapies (e. g. , prophylactic or therapeutic agents) to "manage" a disease so as to prevent the progression or worsening of the disease.
- 25 The term "TLR3 agonist" refers to an affinity agent (i.e., a molecule that binds a target molecule) capable of activating a TLR3 polypeptide to induce a full or partial receptor-mediated response. For example, an agonist of TLR3 induces TLR3-mediated signalling, either directly or indirectly. A TLR3 agonist, as used herein may, but is not required to, bind a TLR3 polypeptide, and may or may not interact directly with the TLR3 polypeptide.
- 30 A "nucleotide agonist" or "nucleic acid agonist" refers to the situation where the affinity agent comprises or consists of nucleotides and/or nucleic acid(s). An "antibody agonist" refers to the situation where the affinity agent is an antibody.
- The terms "polynucleotide" and "nucleic acid", used interchangeably herein, refer to polymeric forms of nucleotides of any length, either ribonucleotides or deoxynucleotides. Thus, these terms
- 35 include, but are not limited to, single-, double-, or multi-stranded DNA or RNA, genomic DNA, cDNA, DNA-RNA hybrids, or a polymer comprising purine and pyrimidine bases or other natural, chemically or biochemically modified, non-natural, or derivatized nucleotide bases. These terms

further include, but are not limited to, mRNA or cDNA that comprise intronic sequences (see, e.g., Niwa et al. (1999) Cell 99(7):691-702). The backbone of the polynucleotide can comprise sugars and phosphate groups (as may typically be found in RNA or DNA), or modified or substituted sugar or phosphate groups. Alternatively, the backbone of the polynucleotide can comprise a
5 polymer of synthetic subunits such as phosphoramidates and thus can be an oligodeoxynucleoside phosphoramidate or a mixed phosphoramidate-phosphodiester oligomer. Peyrottes et al. (1996) Nucl. Acids Res. 24:1841-1848; Chaturvedi et al. (1996) Nucl. Acids Res. 24:2318-2323. A polynucleotide may comprise modified nucleotides, such as methylated nucleotides and nucleotide analogs, uracil, other sugars, and linking groups such as fluororibose and thioate, and nucleotide
10 branches. The sequence of nucleotides may be interrupted by non-nucleotide components. A polynucleotide may be further modified after polymerization, such as by conjugation with a labelling component. Other types of modifications included in this definition are caps, substitution of one or more of the naturally occurring nucleotides with an analog, and introduction of means for attaching the polynucleotide to proteins, metal ions, labelling components, other polynucleotide, or
15 a solid support.

As used herein, the term "host cell" includes a particular subject cell transfected with a nucleic acid molecule and the progeny or potential progeny of such a cell.

The percent identity between the two sequences is a function of the number of identical positions shared by the sequences (i.e., % identity = number of identical overlapping positions/total number of
20 positions x 100%). In one embodiment, the two sequences are the same length. The determination of percent identity between two sequences can also be accomplished using a mathematical algorithm. A preferred, non-limiting example of a mathematical algorithm utilized for the comparison of two sequences is the algorithm of Karlin and Altschul, 1990, Proc. Natl. Acad. Sci. U.S.A. 87:2264-2268, modified as in Karlin and Altschul, 1993, Proc. Natl. Acad. Sci. U.S.A.
25 90:5873. Such an algorithm is incorporated into the NBLAST and XBLAST programs of Altschul et al., 1990, J. Mol. Biol. 215:403. BLAST nucleotide searches can be performed with the NBLAST nucleotide program parameters set, e.g., for score=100, wordlength=12 to obtain nucleotide sequences used. The percent identity between two sequences can be determined using techniques similar to those described above, with or without allowing gaps. In calculating percent
30 identity, typically only exact matches are counted.

As used herein, the term "hybridizes under stringent conditions" describes conditions for hybridization and washing under which nucleotide sequences at least 30% (preferably, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or 98%) identical to each other typically remain hybridized to each other. Such stringent conditions are known to those skilled in
35 the art and can be found in Current Protocols in Molecular Biology, John Wiley & Sons, N.Y. (1989). In one, non-limiting example stringent hybridization conditions are hybridization at 6X sodium chloride/sodium citrate ("SSC") at about 45°C, followed by one or more washes in 0.1X

SSC, 0.2% SDS at about 68°C. In a preferred, non-limiting example stringent hybridization conditions are hybridization in 6X SSC at about 45°C, followed by one or more washes in 0.2X SSC, 0.1% SDS at 50-65°C (i.e., one or more washes at 50°C, 55°C, 60°C or 65°C). It is understood that the nucleic acids of the invention do not include nucleic acid molecules that hybridize under these conditions solely to a nucleotide sequence consisting of only A or T nucleotides. In a particular embodiment, typical stringent hybridisation conditions include temperatures above 30°C, preferably above 35°C, more preferably in excess of 42°C, and/or salinity of less than about 500 mM, preferably less than 200 mM. Hybridization conditions may be adjusted by the skilled person by modifying the temperature, salinity and/or the concentration of other reagents such as SDS, SSC, etc.

TLR3

"TLR3", "TLR3 protein" and "TLR3 receptor", used interchangeably, are used herein to refer to Toll Like Receptor 3, a member of the Toll-like receptor (TLRs) family. Its amino acid sequence of is shown in SEQ ID NO 2 (NCBI accession number NP_003256, the disclosure of which is incorporated herein by reference). Toll Like Receptor 3 is a member of the Toll-like receptor (TLR) family which plays a fundamental role in pathogen recognition and activation of innate immunity. TLRs are highly conserved from Drosophila to humans and share structural and functional similarities. They recognize pathogen-associated molecular patterns (PAMPs) that are expressed on infectious agents, and mediate the production of cytokines necessary for the development of effective immunity. The various TLRs exhibit different patterns of expression. This receptor is most abundantly expressed in placenta and pancreas, and is restricted to the dendritic subpopulation of the leukocytes. It recognizes dsRNA associated with viral infection, and induces the activation of NF-κB and the production of type I interferons. It may thus play a role in host defense against viruses. TLR3 mRNA sequence is described in NCBI accession number NM_003265, the sequence of which is shown in SEQ ED No 1. TLR3 is described in WO 98/50547 (the disclosure of which is incorporated herein by reference).

As used in the present application, the term "TLR3 gene" designates the Toll Like Receptor 3 gene, as well as variants, analogs and fragments thereof, including alleles thereof (e.g., germline mutations). Such variants include, for instance, naturally-occurring variants due to allelic variations between individuals (e.g., polymorphisms), alternative splicing forms, etc. Variants are preferably substantially homologous to NM_003265 sequence, i.e., exhibit a nucleotide sequence identity of at least about 65%, typically at least about 75%, preferably at least about 85%, more preferably at least about 95% with NM 003265 sequence. A particular example of a TLR3 gene comprises NM 003265 sequence. Variants and analogs of a TLR3 gene also include nucleic acid sequences, which hybridize to a sequence as defined above (or a complementary strand thereof) under stringent hybridization conditions. Genetic polymorphisms of the human TLR3 DNA sequence are

known, for example allelic variations in the cytoplasmic region of TLR3 gene and in the immediate 5' sequence of the TLR3 gene (see Pirieli et al. (2005) Tissue Antigens 66(2): 125, the disclosure of which is incorporated herein by reference), for example the C/T polymorphism at position 2593, the C/A polymorphism at position 2642 and the *A/G* polymorphism at position 2690 in the TLR3 gene.

A fragment of a TLR3 gene designates any portion of at least about 8 consecutive nucleotides of a sequence as disclosed above, preferably at least about 15, more preferably at least about 20 nucleotides, further preferably of at least 30 nucleotides. Fragments include all possible nucleotide lengths between 8 and 100 nucleotides, preferably between 15 and 100, more preferably between 20 and 100.

The term "gene" shall be construed to include any type of coding nucleic acid, including genomic DNA (gDNA), complementary DNA (cDNA), synthetic or semi-synthetic DNA, as well as any form of corresponding RNA. The term gene particularly includes recombinant nucleic acids encoding TLR3, i.e., any non naturally occurring nucleic acid molecule created artificially, e.g., by assembling, cutting, ligating or amplifying sequences. A TLR3 gene is typically double-stranded, although other forms may be contemplated, such as single-stranded. TLR3 genes may be obtained from various sources and according to various techniques known in the art, such as by screening DNA libraries or by amplification from various natural sources. Recombinant nucleic acids may be prepared by conventional techniques, including chemical synthesis, genetic engineering, enzymatic techniques, or a combination thereof.

A "TLR3 polypeptide" designates any molecule comprising a stretch of at least six consecutive amino acids present in a TLR3 protein. A TLR3 polypeptide may be encoded by a fragment of a TLR3 gene, may be synthetically produced or may be produced through proteolytic digestion of a TLR3 protein or another TLR3 polypeptide. A TLR3 polypeptide may be modified, such as by glycosylations and/or acetylations and/or chemical reaction or coupling, and may contain one or several non-natural or synthetic amino acids. A specific example of a TLR3 polypeptide comprises all or part of NP_003256 sequence.

TLR3 agonists

1. Assaying TLR3 Agonist Activity

A TLR3 agonist according to the present invention can be selected from any suitable agent that activates TLR3 and/or the subsequent cascade of biochemical events associated with TLR3 activation in vivo. Assays for detecting TLR3 agonist activity are known in the art and include, for example, the detection of luciferase (luc) production from an NF- κ B reporter plasmid, or the induction of endogenous IL-8 (K. Kariko et al., *J. Immunol.* 2004,172: 6545-49, the disclosures of which is incorporated herein by reference). Assays for detecting TLR3 agonism of test compounds are also described, for example, in PCT publication Nos. WO 03/31573, WO 04/053057, WO

04/053452, and WO 04/09467 1, the disclosures of each of which are incorporated herein by reference.

Regardless of the particular assay employed, a compound can be identified as an agonist of TLR3 if performing the assay with that compound results in at least a threshold increase of some biological activity known to be mediated by TLR3. Conversely, a compound may be identified as not acting as an agonist of TLR3 if, when used to perform an assay designed to detect biological activity mediated by TLR3, the compound fails to elicit a threshold increase in the biological activity. Unless otherwise indicated, an increase in biological activity refers to an increase in the same biological activity over that observed in an appropriate control. An assay may or may not be performed in conjunction with the appropriate control. With experience, one skilled in the art may develop sufficient familiarity with a particular assay (e.g., the range of values observed in an appropriate control under specific assay conditions) that performing a control may not always be necessary to determine the TLR3 agonism of a compound in a particular assay. The precise threshold increase of TLR3-mediated biological activity for determining whether a particular compound is or is not an agonist of TLR3 in a given assay may vary according to factors known in the art including but not limited to the biological activity observed as the endpoint of the assay, the method used to measure or detect the endpoint of the assay, the signal-to-noise ratio of the assay, the precision of the assay, and whether the same assay is being used to determine the agonism of a compound for multiple TLRs. Accordingly it is not practical to set forth generally the threshold increase of TLR3-mediated biological activity required to identify a compound as being an agonist or a non-agonist of TLR3 for all possible assays. Those of ordinary skill in the art, however, can readily determine the appropriate threshold with due consideration of such factors. However, regardless of the particular assay employed, a compound can generally be identified as an agonist of TLR3 if performing the assay with a compound results in at least a threshold increase of some biological activity mediated by TLR3.

Assays employing HEK293 cells transfected with an expressible TLR3 structural gene may use a threshold of, for example, at least a three-fold increase in a TLR3-mediated biological activity (e.g., NF-KB activation) when the compound is provided at a concentration of, for example, from about 1 μ M to about 10 μ M for identifying a compound as an agonist of the TLR3 transfected into the cell. However, different thresholds and/or different concentration ranges may be suitable in certain circumstances. Also, different thresholds may be appropriate for different assays.

2. TLR3 Agonistic Antibodies

A TLR3 agonist of this invention may be an agonistic antibody, an agonistic fragment of such antibodies, a chimeric version of such antibodies or fragment, or another active antibody derivative. TLR3 agonist antibodies useful in this invention may be produced by any of a variety of techniques known in the art. Typically, they are produced by immunization of a non-human animal, preferably

- a mouse, with an immunogen comprising a TLR3 protein or TLR3 peptide. The immunogen may comprise intact TLR3-expressing tumor cells, cell membranes from TLR-3-expressing cells, the full length sequence of the TLR3 protein (produced recombinantly or isolated from a natural source), or a fragment or derivative thereof, typically an immunogenic fragment, i.e., a portion of the polypeptide comprising an epitope exposed on the surface of cells expressing the TLR3. An immunogenic fragment typically contains at least 7 consecutive amino acids present in the full-length TLR3 amino acid sequence, even more preferably at least 10 consecutive amino acids thereof. The amino acid sequence of the fragment is essentially derived from the extracellular domain of the TLR3 protein. In preferred embodiments, the TLR3 protein or peptide used to generate antibodies is a human TLR3 protein or peptide.
- In a most preferred embodiment, the immunogen comprises a wild-type human TLR3 polypeptide in a lipid membrane, typically derived from a membrane fraction of a TLR-3 expressing cell. In a specific embodiment, the immunogen comprises whole TLR3 expressing tumor cells, intact or optionally chemically or physically lysed.
- The preparation of monoclonal or polyclonal antibodies is well known in the art, and any of a large number of available techniques can be used (see, e.g., Kohler & Milstein, *Nature* 256:495-497 (1975); Kozbor et al., *Immunology Today* 4: 72 (1983); Cole et al., pp. 77-96 in *Monoclonal Antibodies and Cancer Therapy* (1985)). The step of immunizing a non-human mammal with an immunogen may be carried out in any manner well known in the art for (see, for example, E. Harlow and D. Lane, Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY (1988)). Generally, the immunogen is suspended or dissolved in a buffer, optionally with an adjuvant, such as complete Freund's adjuvant. Methods for determining the amount of immunogen, types of buffers and amounts of adjuvant are well known to those of skill in the art and are not limiting in any way on the present invention.
- Similarly, the location and frequency of immunization sufficient to stimulate the production of antibodies is also well known in the art. In a typical immunization protocol, the non-human animals are injected intraperitoneally with antigen on day 1 and again about a week later. This is followed by recall injections of the antigen around day 20, optionally with adjuvant such as incomplete Freund's adjuvant. The recall injections are performed intravenously and may be repeated for several consecutive days. This is followed by a booster injection at day 40, either intravenously or intraperitoneally, typically without adjuvant. This protocol results in the production of antigen-specific antibody-producing B cells after about 40 days. Other protocols may also be utilized as long as they result in the production of B cells expressing an antibody directed to the antigen used in immunization.
- In another embodiment, lymphocytes from an unimmunized non-human mammal are isolated, grown in vitro, and then exposed to the immunogen in cell culture. The lymphocytes are then harvested and the fusion step described below is carried out.

For monoclonal antibodies, which are preferred for producing agonistic antibodies for use in the present invention, the next step is the isolation of cells, e.g., lymphocytes, splenocytes, or B cells, from the immunized non-human mammal and the subsequent fusion of those splenocytes, or B cells, or lymphocytes, with an immortalized cell in order to form an antibody-producing

5 hybridoma. The isolation of splenocytes, e.g., from a non-human mammal is well-known in the art and, e.g., involves removing the spleen from an anesthetized non-human mammal, cutting it into small pieces and squeezing the splenocytes from the splenic capsule and through a nylon mesh of a cell strainer into an appropriate buffer so as to produce a single cell suspension. The cells are washed, centrifuged and resuspended in a buffer that lyses any red blood cells. The solution is

10 again centrifuged and remaining lymphocytes in the pellet are finally resuspended in fresh buffer. Once isolated and present in single cell suspension, the antibody-producing cells are fused to an immortal cell line. This is typically a mouse myeloma cell line, although many other immortal cell lines useful for creating hybridomas are known in the art. Preferred murine myeloma lines include, but are not limited to, those derived from MOPC-21 and MPC-1 1 mouse tumors available from the

15 SaIk Institute Cell Distribution Center, San Diego, Calif. U.S.A., X63 Ag8653 and SP-2 cells available from the American Type Culture Collection, Rockville, Maryland U.S.A. The fusion is effected using polyethylene glycol or the like. The resulting hybridomas are then grown in selective media that contains one or more substances that inhibit the growth or survival of the unfused, parental myeloma cells. For example, if the parental myeloma cells lack the enzyme hypoxanthine

20 guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine (HAT medium), which substances prevent the growth of HGPRT-deficient cells.

The hybridomas can be grown on a feeder layer of macrophages. The macrophages are preferably from littermates of the non-human mammal used to isolate splenocytes and are typically primed

25 with incomplete Freund's adjuvant or the like several days before plating the hybridomas. Fusion methods are described, e.g., in (Goding, "Monoclonal Antibodies: Principles and Practice," pp. 59-103 (Academic Press, 1986)), the disclosure of which is herein incorporated by reference.

The cells are allowed to grow in the selection media for sufficient time for colony formation and antibody production. This is usually between 7 and 14 days. The supernatants from the hybridoma

30 colonies are then assayed for the production of antibodies that specifically activate the TLR3 protein. The wells positive for the desired antibody production are examined to determine if one or more distinct colonies are present. If more than one colony is present, the cells may be re-cloned and grown to ensure that only a single cell has given rise to the colony producing the desired antibody. Positive wells with a single apparent colony are typically recloned and re-assayed to

35 ensure that only one monoclonal antibody is being detected and produced.

Hybridomas that are confirmed to be producing a TLR3 agonistic monoclonal antibody are then grown up in larger amounts in an appropriate medium, such as DMEM or RPMI-1640.

Alternatively, the hybridoma cells can be grown in vivo as ascites tumors in an animal.

After sufficient growth to produce the desired monoclonal antibody, the growth media containing
5 monoclonal antibody (or the ascites fluid) is separated away from the cells and the monoclonal antibody present therein is purified. Purification is typically achieved by gel electrophoresis, dialysis, chromatography using protein A or protein G-Sepharose, or an anti-mouse Ig linked to a solid support such as agarose or Sepharose beads (all described, for example, in the Antibody Purification Handbook, Amersham Biosciences, publication No. 18-1037-46, Edition AC, the
10 disclosure of which is hereby incorporated by reference). The bound antibody is typically eluted from protein A/protein G columns by using low pH buffers (glycine or acetate buffers of pH 3.0 or less) with immediate neutralization of antibody-containing fractions. These fractions are pooled, dialyzed, and concentrated as needed.

In certain embodiments, the DNA encoding an antibody that agonizes TLR3 is isolated from the
15 hybridoma, and placed in an appropriate expression vector for transfection into an appropriate host. The host is then used for the recombinant production of the antibody, variants thereof, active fragments thereof, or humanized or chimeric antibodies comprising the antigen recognition portion of the antibody.

DNA encoding the monoclonal antibodies of the invention can be readily isolated and sequenced
20 using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies). Once isolated, the DNA can be placed into expression vectors, which are then transfected into host cells such as E. coli cells, simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of monoclonal antibodies in the
25 recombinant host cells. Recombinant expression in bacteria of DNA encoding the antibody is well known in the art (see, for example, Skerra et al. (1993) Curr. Op. Immunol. 5:256; and Pluckthun (1992) Immunol. Revs. 130:151. Antibodies may also be produced by selection of combinatorial libraries of immunoglobulins, as disclosed for instance in Ward et al. (1989) Nature 341:544.

The TLR3 agonist antibodies can be full length antibodies or antibody fragments or derivatives.

30 Examples of antibody fragments include Fab, Fab', Fab'-SH, F(ab')₂, and Fv fragments; diabodies; single-chain Fv (scFv) molecules; single chain polypeptides containing only one light chain variable domain, or a fragment thereof that contains the three CDRs of the light chain variable domain, without an associated heavy chain moiety; single chain polypeptides containing only one heavy chain variable region, or a fragment thereof containing the three CDRs of the heavy chain
35 variable region, without an associated light chain moiety; and multispecific antibodies formed from antibody fragments. Such fragments and derivatives and methods of preparing them are well known in the art. For example, pepsin can be used to digest an antibody below the disulfide

linkages in the hinge region to produce F(ab)'₂, a dimer of Fab which itself is a light chain joined to VH-CH1 by a disulfide bond. The F(ab)'₂ may be reduced under mild conditions to break the disulfide linkage in the hinge region, thereby converting the F(ab)'₂ dimer into an Fab' monomer. The Fab' monomer is essentially Fab with part of the hinge region (see Fundamental Immunology (Paul ed., 3d ed. 1993)). While various antibody fragments are defined in terms of the digestion of an intact antibody, one of skill will appreciate that such fragments may be synthesized de novo either chemically or by using recombinant DNA methodology.

3. Small Molecule TLR3 Agonists

A small molecule TLR3 agonist is preferably an organic molecule of less than about 1500 daltons. The design and selection (e.g., from a combinatorial library) or synthesis of a small molecule TLR3 agonist may be achieved through the use of the known crystal structure of TLR3 (Choe et al., Science, 309, pp. 581-85 (2005), the disclosure of which is herein incorporated by reference). This design or selection may begin with selection of the various moieties which fill the putative binding pocket(s) in which known double-stranded RNA agonists bind. There are a number of ways to select moieties to fill individual binding pockets. These include visual inspection of a physical model or computer model of the active site elucidated from the crystal structure and manual docking of models of selected moieties into various binding pockets. Modelling software that is well known and available in the art may be used. These include QUANTA [Molecular Simulations, Inc., Burlington, Mass., 1992], and SYBYL [Molecular Modelling Software, Tripos Associates, Inc., St. Louis, Mo., 1992]. This modelling step may be followed by energy minimization with standard molecular mechanics force fields such as CHARMM and AMBER. [AMBER: (S. J. Weiner, P. A. Kollman, D. A. Case, U. C. Singh, C. Ghio, G. Alagona, and P. Weiner, J. Am. Chem. Soc., 1984, 106, 765); CHARMM: (B. R. Brooks, R. E. Bruccoleri, B. D. Olafson, D. J. States, S Swaminathan, and M. Karplus, J. Comp. Chem. 1983, 4, 187). In addition, there are a number of more specialized computer programs to assist in the process of optimally placing either complete molecules or molecular fragments into the TLR3 agonist binding site. These include: GRID (Goodford, P. J. A Computational Procedure for Determining Energetically Favorable Binding Sites on Biologically Important Macromolecules. J. Med. Chem. 1985, 28, 849-857). GRID is available from Oxford University, Oxford, UK; MCSS (Mariner, A.; Karplus, M. Functionality Maps of Binding Sites: A Multiple Copy Simultaneous Search Method. Proteins: Structure, Function and Genetics 1991, 11, 29-34). MCSS is available from Molecular Simulations, Burlington, Mass.; and DOCK (Kuntz, I. D.; Blaney, J. M.; Oatley, S. J.; Langridge, R.; Ferrin, T. E. A Geometric Approach to Macromolecule-Ligand Interactions. J. Mol. Biol. 1982, 161, 269-288). DOCK is available from the University of California, San Francisco, Calif.

Once suitable binding orientations have been selected, complete molecules can be chosen for biological evaluation. In the case of molecular fragments, they can be assembled into a single agonist. This assembly may be accomplished by connecting the various moieties to a central scaffold. The assembly process may, for example, be done by visual inspection followed by manual
5 model building, again using software such as Quanta or Sybyl. A number of other programs may also be used to help select ways to connect the various moieties. These include: CAVEAT (Bartlett, P. A.; Shea, G. T.; Telfer, S. J.; Waterman, S. CAVEAT: A Program to Facilitate the Structure-Derived Design of Biologically Active Molecules. In "Molecular Recognition in Chemical and Biological Problems," Special Pub., Royal Chem. Soc. 1989, 78, 182-196). CAVEAT is available
10 from the University of California, Berkeley, Calif.; 3D Database systems, such as MACCS-3D (MDL Information Systems, San Leandro, Calif.). This area has been reviewed by Martin (Martin, Y. C. 3D Database Searching in Drug Design. J. Med. Chem. 1992, 35, 2145); and HOOK (available from Molecular Simulations, Burlington, Mass.).

In addition to the above computer assisted modelling of agonist compounds, a TLR3 agonist may
15 be constructed "de novo" using either an empty agonist binding site of TLR3 or optionally including some portions of a known agonist. Such methods are well known in the art. They include, for example: LUDI (Bohm, H. J. The Computer Program LUDI: A New Method for the De Novo Design of Enzyme Inhibitors. J. Comp. Aid. Molec. Design. 1992, 6, 61-78). LUDI is available from Biosym Technologies, San Diego, Calif.; LEGEND (Nishibata, Y., Itai, A., Tetrahedron,
20 1991, 47, 8985). LEGEND is available from Molecular Simulations, Burlington, Mass.; and LeapFrog (available from Tripos Associates, St. Louis, Mo.).

A number of techniques commonly used for modelling drugs may be employed (For a review, see: Cohen, N. C ; Blaney, J. M.; Humblet, C ; Gund, P.; Barry, D. C , "Molecular Modeling Software and Methods for Medicinal Chemistry", J. Med. Chem., 1990, 33, 883). There are likewise a
25 number of examples in the chemical literature of techniques that can be applied to specific drug design projects. For a review, see: Navia, M. A. and Murcko, M. A., Current Opinions in Structural Biology. 1992, 2, 202. Some examples of these specific applications include: Baldwin, J. J. et al., J. Med. Chem., 1989, 32, 2510; Appelt, K. et al., J. Med. Chem., 1991, 34, 1925; and Ealick, S. E. et al., Proc. Nat. Acad. Sci. USA. 1991, 88, 11540.

30 An alternative to designing or modelling small molecule TLR3 agonists is to screen existing small molecule chemical libraries for a TLR3 agonist. Such libraries can be screened by any TLR3 agonist assay known in the art, including those described herein. The compound libraries may be initially screened using a higher throughput assay, such as a competition assay with a known, labelled TLR3 agonist, such as the double-stranded RNA molecules polyI:polyC or polyA:polyU.
35 Compounds that are positive in a competition assay are then further assayed for their ability to activate TLR3 and cause the subsequent cascade of biochemical events.

Thus, according to one embodiment, the invention provides a kit for determining if a test compound is useful for the treatment of cancer comprising a cancer cell characterized by the presence of TLR3 on its cell surface; and a reagent for detecting apoptosis or cell death. Such a kit optionally comprises a reagent that detects agonism of TLR3 by said test compound. This optional
5 component may be used to confirm that any apoptosis or cell death of the cancer cell caused by said test compound is TLR3-mediated.

4. Nucleic Acid-based TLR3 Agonists

Nucleic acid-based TLR3 agonists useful in this invention comprise a region of double-stranded
10 ribonucleic acids. The term "double-stranded" means a portion of the agonist where ribonucleotides are hydrogen bonded (base-paired) to complementary ribonucleotides to form a double-stranded structure. Preferably the entire nucleic acid-based TLR3 agonist consists of ribonucleotides and chemically modified ribonucleotides ("dsRNA TLR3 agonist"). More preferably, at least 50% of the dsRNA TLR3 agonist is in a double-stranded conformation under in
15 vivo conditions. Even more preferred is if at least 60%, 70%, 80%, 90%, 95%, or 97% of the dsRNA TLR3 agonist is in a double-stranded conformation under in vivo conditions. The determination of what percentage of the dsRNA TLR3 agonist is in a double-stranded conformation is achieved by dividing the number of nucleotides that are base-paired by the total number of nucleotides in a molecule. Thus, a 21 base-paired molecule containing 2 nucleotide
20 overhangs at both the 3' and 5' end would have 42 nucleotides that are base-paired and 4 nucleotides that are not base-paired, making it 42/46 or 91.3% double-stranded. Similarly, a molecule comprised of two 21 nucleotide strands that are complementary to one another at all nucleotides except for two nucleotides within the middle of each strand would have 38 (19 +19)
25 nucleotides that were base-paired and 4 (2+2) that were not base-paired. Such a molecule would be 38/42 or 90.5% double-stranded.

A double-stranded region of a dsRNA TLR3 agonist can be formed by a self-complementary region of a single RNA molecule (e.g., a stem and loop structure, such as hairpin RNA and shRNA), by two molecules of RNA that hybridize with one another in whole or in part (as in double-stranded RNA), or a mixture of both (e.g., a partially self-complementary molecule of RNA
30 and a second RNA molecule that hybridizes to regions in the former in that remain single-stranded after the formation of the hairpin). The dsRNA TLR3 agonist of this invention may also comprise single-stranded regions, such as 3' and/or 5' overhangs at either end of the agonist, and/or "mismatched" or "loop-out" structures within the agonist.

In the case of shRNA, the dsRNA TLR3 agonist is encoded by a DNA sequence present on an
35 expression vector. The expression vector is the molecule that is administered to the subject. The expression vector typically comprises a promoter that is activated by RNA polymerase II or III and terminator sequences, each of which is operably linked to the shRNA coding sequence to ensure its

proper transcription. Promoters activated by RNA polymerase π include, but are not limited to, U6, tRNA^{Val}, H1, and modified versions of the foregoing. Promoters activated by RNA polymerase III include, but are not limited to, CMV and EF1 α . In one preferred embodiment, the promoter is an inducible promoter or a tumor cell-specific promoter.

- 5 Within the context of the present invention, the term "dsRNA TLR3 agonist" designates any therapeutically or prophylactically effective RNA compound that comprises a double-stranded region. Such compounds are typically active per se, i.e., they do not encode a polypeptide or do not require translation to be active. A dsRNA TLR3 agonist can be of any length. Preferably, a dsRNA TLR3 agonist has a length of at least about 10 base pairs (bp), 20bp, 30bp, 50bp, 80bp, 100bp, 200bp, 400bp, 600bp, 800bp or 1000bp. In one aspect the dsRNA TLR3 agonist is a short dsRNA having a chain length of less than 30bp, 50bp, 80bp, 100bp or 200bp. In another embodiment, the dsRNA TLR3 agonist is a longer dsRNA, but having a chain length of less than 400bp, 600bp, 800bp or 1000bp. In another embodiment, the dsRNA TLR3 agonist is a long dsRNA having a chain length of greater than 1000bp.
- 10 15 In one aspect, a dsRNA TLR3 agonist is a composition that comprises a heterogeneous mixture of dsRNA molecules, wherein a plurality of molecules have differing lengths. Preferably the dsRNA molecules in such a composition have on average a length of at least about 10bp, 20bp, 30bp, 50bp, 80bp, 100bp, 200bp, 400bp, 600bp, 800bp or 1000bp. In another embodiment, a dsRNA TLR3 agonist composition comprises a plurality dsRNA molecules where at least 20%, 50%, 80%, 90% or 98% of dsRNA molecules have a length of at least about 10bp, 20bp, 30bp, 50bp, 80bp, 100bp, 200bp, 400bp, 600bp, 800bp or 1000bp per strand. In one preferred example, the dsRNA is a short dsRNA having between 10 and 30, more preferably between 20 and 30bp per strand. In a preferred embodiment a dsRNA TLR3 agonist composition has a substantially homogenous mixture of dsRNA molecules, where substantially all the molecules on each strand do not differ in chain length by more than 30bp, 50bp, 80bp, 100bp or 200bp. Average chain length of dsRNA TLR3 agonists in a composition can be determined easily, for example, by gel permeation chromatography. One or more of the dsRNA molecules within such a compositions is optionally a siRNA molecule targeted against a cancer antigen.
- 20 25 30 35 The ribonucleotides in the dsRNA TLR3 agonist of this invention can be natural or synthetic, and may be chemically modified derivatives or analogs of natural nucleotides. Preferred modification are stabilizing modifications, and thus can include at least one modification in the phosphodiester linkage and/or on the sugar, and/or on the base. For example, one or both strands of the dsRNA can independently include one or more phosphorothioate linkages, phosphorodithioate linkages, and/or methylphosphonate linkages; modifications at the 2'-position of the sugar, such as 2'-O-methyl modifications, 2'-O-methoxyethyl modifications, 2'-amino modifications, 2'-deoxy modifications, 2'-halo modifications such as 2'-fluoro; combinations of the above, such as 2'-deoxy-2'-fluoro modifications; acyclic nucleotide analogs, and can also include at least one phosphodiester linkage.

Oligonucleotides used in the dsRNA TLR3 agonist of this invention may also include base modifications or substitutions. Modified bases include other synthetic and naturally-occurring bases such as 5-methylcytosine (5-Me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine and inosine, 2-propyl and other
5 alkyl derivatives of adenine and inosine, 2-thiouracil and 2-thiocytosine, 5-halouracil and cytosine, 5-propynyl(-C≡C-CH₃) uracil and cytosine and other alkynyl derivatives of pyrimidine bases, 6-azouracil and cytosine, 5-uracil (pseudouracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, S-thioalkyl, 8-hydroxyl and other 8-substituted adenines and inosines, 5-halo particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylinosine and 7-methyladenine, 2-F-adenine, 2-amino-adenine, 8-azainosine and 8-azaadenine, 7-deazainosine and 7-deazaadenine and 3-deazainosine and 3-deazaadenine.

Other modifications include a 3'- and/or 5'-terminal cap, a terminal 3'-5' linkage, and a 5'-terminal phosphate group or modified phosphate group. Examples of terminal cap moieties include, but are not limited to, an inverted deoxy abasic moiety, an inverted deoxynucleotides, or a glyceryl moiety.
15 In addition one or both strands (in a two-stranded dsRNA TLR3 agonist) may be or include a concatemer consisting of two or more oligonucleotide sequences joined by a linker(s). All of these modifications are well known in the art (see, for example, Kandimalla et al. ((2003) Nucl. Acid. Res. 31(9): 2393-2400).

Previous studies of double-stranded RNA (dsRNA) assessing their ability to be effective interferon
20 inducers suggested that dsRNA agents must possess the secondary structure of a double stranded helix. Other dsRNA agents which have also been shown to be suitable as TLR3 agonist include double-stranded polynucleotides which are not complementary or not perfectly complementary; these have been known as, so-called "mismatched" or "loop-out" structures and exist in naturally occurring RNAs such as transfer tRNAs, ribosomal RNAs and the viral RNA secondary structures.
25 One commonly cited dsRNA compound, Ampligen, comprises a structure where a few parts of cytidine in the poly Lpoly C structure are replaced with uridine (i.e. mismatched RNA); this compound has been reported to have physiological activity similar to that of the parent poly lipoly C. However it will be appreciated that TLR3 agonists of any type and configuration can be used in accordance with this invention.

30 Generally, the polynucleotides need to be resistant to nucleases in order to remain as macromolecules for a sufficient length of time; polynucleotides are less sensitive to nuclease attack when they are in a helical complex. However, certain analogs such as AmpligenTM appear to retain their TLR3 agonist activity.

In a particular embodiment, each strand of these dsRNAs can have a length comprised between
35 about 5 and 50 bases, more preferably between 5 and 40, 35, 30, 25 or 20 bases. Each strand is preferably perfectly complementary to the other. Preferred examples of such dsRNAs are

homopolyRNAs, i.e., dsRNAs in which each strand comprises essentially a repeat of the same base; or comprises a homopolyRNA region.

The base in such homopolyRNA strands may be any naturally occurring base (e.g., polyA, polyU, polyC, polyG) or non-naturally occurring (e.g., chemically synthesized or modified) base (e.g.,

5 polyI). Polynucleotides typified by polyinosinic-polycytidylic acid, i.e., poly(I):poly(C) or poly LC and polyadenylic-polyuridylic acid, i.e., poly(A):poly(U) or poly A:U, are well-known compounds in the art and have been known to induce interferon production by immune cells. Thus in preferred embodiments, the TLR3 agonist for use according to the invention is a double stranded RNA selected from the group consisting of: polyinosinic acid and polycytidylic acid, polyadenylic acid
10 and polyuridylic acid, polyinosinic acid analogue and polycytidylic acid, polyinosinic acid and polycytidylic acid analogue, polyinosinic acid analogue and polycytidylic acid analogue, polyadenylic acid analogue and polyuridylic acid, polyadenylic acid and polyuridylic acid analogue, and polyadenylic acid analogue and polyuridylic acid analogue. The term "analogue" as used herein means any of the nucleotide modifications described above.

15 It will be appreciated that dsRNA TLR3 agonists can comprise any combination of bases and be designed using any suitable method. Preferably, the basic requirement of a region of double-strandedness, stability and resistance to nuclease attack and the preferences for chain length are taken into account. These properties, as well as relative TLR3 agonistic activity of any dsRNA TLR3 agonist can be tested and assessed with reference to the a $rA_n:rU_n$ or $rI_n:rC_n$ complex for
20 example. Measures can be taken to increase stability and resistance to nucleases, or to increase or optionally decrease interferon-inducing action.

Other examples of dsRNA include nucleic acids described in U.S. Patent Nos. 5,298,614 and 6,780,429. U.S. Patent no. 5,298,614 reports that when chain length of the double stranded nucleic acid derivatives is limited to certain ranges, the resulting substances exhibit desired physiological
25 activity with markedly less toxicity, providing polynucleotides having a length of about 50 to 10,000 as calculated by base pair numbers. Also described are derivative wherein the purine or pyrimidine ring in the nucleic acid polymer is substituted with at least one SH group, or said derivative contains a disulphide bond, or both (preferred ratio of number of sulphur atoms to cytidylic acid present in the poly C are 1:6 to 39). U.S. Patent No. 6,780,429 describes a particular
30 type of dsRNA compounds that are "chain-shortened" having lengths of about 100 to 1,000 as calculated by base pair numbers, or preferably from 200 to 800, and more preferably from 300 to 600. The latter compounds are reported to contain low numbers of 2'-5' phosphodiester bonds by a method designed to avoid phosphate groups causing intramolecular rearrangement from 3' position to 2' position through a mechanism called pseudo rotation simultaneously that can occur during
35 hydrolysis of polynucleotides, resulting in a portion of 3'-5' phosphodiester bonds in the chain-shortened polynucleotide molecule being replaced by 2'-5' phosphodiester bonds. The disclosures of each of these references is incorporated herein by reference.

Other nucleic acid agonists that can be suitable for use as TLR3 agonists are provided in: Field et al: Proc. Nat. Acad. Sci. U.S. 58, 1004, (1967); Field et al: Proc. Nat. Acad. Sci. U.S. 58, 2102, (1967); Field et al: Proc. Nat. Acad. Sci. U.S. 61, 340, (1968); Tytell et al: Proc. Nat. Acad. Sci. U.S. 58, 1719, (1967); Field et al: J. Gen. Physiol. 56, 905 (1970); De Clercq et al: Methods in Enzymology, 78, 291 (1981). A number of synthetic nucleic acid derivatives have been described, including homopolymer-homopolymer complexes (Double Strand Nucleic Acid Polymer such as those in which poly I:C or poly A:U are a parent structure, where these homopolymer-homopolymer complexes contain: (1) base modifications, exemplified by Polyinosinic acid-poly(5-bromocytidylic acid), Polyinosinic acid-poly(2-thiocytidylic acid), Poly(7-deazainosinic acid)-polycytidylic acid, Poly(7-deazainosinic acid)-poly(5-bromocytidylic acid), and Polyinosinic acid-poly(5-thiouridylic acid); (2) Sugar Modifications, exemplified by Poly(2'-azidoinosinic acid)-polycytidylic acid; and (3) Phosphoric Acid Modifications, exemplified by Polyinosinic acid-poly(cytidyl-5'-thiophosphoric acid). Other synthetic nucleic acid derivatives that have been described include interchanged copolymers, exemplified by Poly(adenylic acid-uridylic acid); and homopolymer-copolymer complexes, exemplified by Polyinosinic acid-poly(cytidylic acid-uridylic acid) and Polyinosinic acid-poly(citydylic acid-4-thiouridylic acid). Other synthetic nucleic acid derivatives that have been described include complexes of synthetic nucleic acid with polycation, exemplified by Polyinosinic acid-polycytidylic acid-poly-L-lysinecarboxy-methylcellulose complex (called "Poly ICLC"). Yet another example of synthetic nucleic acid derivative is Polyinosinic acid-poly(l-vinylcytosine).

One example of a TLR3 agonist is Ampligen™ (Hemispherx, Inc., of Rockville, Md., U.S.A.), a dsRNA formed by complexes of polyriboinosinic and polyribocytidylic/uridylic acid, such as $rI_n:r(C_x, U \text{ or } G)_n$ where x has a value from 4 to 29, e.g., $rI_n:r(Cj2 \text{ U})_n$. Many mismatched dsRNA polymers which behave similarly to Ampligen have been studied; mismatched dsRNA based on poly LC have included complexes of a polyinosinate and a polycytidylate containing a proportion of uracil bases or guanidine bases, e.g., from 1 in 5 to 1 in 30 such bases. The key therapeutic advantage of mismatched dsRNAs over other forms of natural and/or synthetic dsRNAs is a reported reduction in toxicity over compounds such as those described in Lampson et al in U.S. Patent No. 3,666,646.

Specific examples of double-stranded RNA according to the present invention further include Polyadenur (Ipsen) and Ampligen (Hemispherx). Polyadenur is a polyA/U RNA molecule, i.e., contains a polyA strand and a polyU strand. Polyadenur has been developed for the potential treatment of hepatitis B virus (HBV) infection. Ampligen is of a polyI/polyC compound (or a variant thereof comprising a polyI/polyC12U RNA molecule). Ampligen is disclosed for instance in EP 281 380 or EP 113 162. Ampligen has been proposed for the treatment of cancer, viral infections and immune disorders. It was developed primarily for the potential treatment of myalgic

encephalomyelitis (ME, or chronic fatigue syndrome/chronic fatigue immune dysfunction syndrome, CFS/CFIDS).

A particular example of a dsRNA for use in the present invention is a dsRNA comprising a polyA/polyU region, wherein each strand of said dsRNA contains less than 25 bases.

- 5 Another particular example of a dsRNA for use in the present invention is a dsRNA comprising a polyI/polyC(U) region, wherein each strand of said dsRNA contains less than 25 bases.

Further dsRNAs have been disclosed in the literature or may be developed, which can be used within the present invention. More generally, any synthetic double-stranded homopolyRNA may be used in the context of this invention, as well as any other dsRNA as herein described.

- 10 In a more preferred embodiment, the dsRNA is a fully double stranded (e.g., blunt-ended; no overhangs) polyA:polyU molecule consisting of between 19 and 30 base pairs (e.g., 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 or 30 base pairs) and comprising between 1 and 30 stabilizing modifications and/or a 3' and/or a 5' cap. Stabilizing modification and caps are described above and are well-known in the art. The stabilizing modifications and/or the presence of a cap make the
- 15 dsRNA more resistant to serum degradation. Preferred stabilizing modifications include phosphorothioate internucleotide linkages, 2'-deoxyribonucleotides, 2'-O-methyl ribonucleotides, 2'-deoxy-2'-fluoro ribonucleotides, "universal base" nucleotides, "acyclic" nucleotides, 5-C-methyl nucleotides, and terminal glyceryl and/or inverted deoxy abasic residue incorporation. These chemical modifications are known to dramatically increase the serum stability of dsRNA
- 20 compounds. One example of a stabilized dsRNA is Stealth™ RNAi (commercially available from Invitrogen, Carlsbad, CA USA).

- In another preferred embodiment the dsRNA TLR3 agonist is a siRNA molecule or a shRNA molecule that is designed to specifically hybridize with the mRNA coding a tumor cell antigen or another protein involved in tumor proliferation. In this embodiment, the dsRNA molecule plays a
- 25 dual role in the treatment of cancer. It is both an agonist of TLR3 and a suppressor of a specific tumor antigen expression. It has been demonstrated that siRNA molecules and shRNA molecules targeted against cellular proteins exhibit both sequence-dependent gene suppression and sequence-independent effects mediated through TLR3 (K. Kariko et al., J. Immunol., 2004, 172: 6545-6549). Thus, it is expected that tumor antigen- or tumor-proliferation specific siRNA and shRNA
- 30 molecules will also be agonists of TLR3 in cancer cells.

- The choice of ribonucleotide sequences to use for such dsRNA TLR3 agonists will depend upon the specific tumor antigen or tumor proliferative protein to be suppressed, as well as the specific type of cancer to be treated. By way of example, the dsRNA TLR3 agonist may have a ribonucleotide sequence designed to hybridize to a mRNA encoding VEGF, VEGF receptors, A-
- 35 Raf, B-Raf, C-Raf, Raf-1, HSP70, HSP90, PDGF, TGF-alpha, EGF, EGF receptor, a member of the human EGF-like receptor family such as HER-2/neu, HER-3, HER-4 or a heterodimeric receptor comprised of at least one HER subunit, carcinoembryonic antigen, gastrin releasing

peptide receptor antigen, Muc-1, CA125, α B3 integrins, α 5B1 integrins, α IIB3-integrins, CTLA-4, CD20, CD22, CD30, CD33, CD52, CD56, CD80, PDGF beta receptor, Src, VE-cadherin, IL-8, hCG, IL-6, *IL-6* receptor, IL-15, or an mRNA encoding any additional protein targets set forth in <http://oncologyknowledgebase.com/oksite/TargetedTherapeutics/TTOExhibit2.pdf> and <http://oncologyknowledgebase.com/oksite/TargetedTherapeutics/TTOExhibit3.pdf> . the disclosures of which are herein incorporated by reference. The gene sequences encoding these proteins are known. Thus, siRNA and shRNA intended to suppress the expression of these proteins can therefore be designed using well-known software and algorithms (see, e.g., <https://rnaidesigner.invitrogen.com/rnaiexpress/>) .

10

Detection of Cancer Cells expressing TLR3

It has also been found that certain classes of patients with cancer treated in accordance with the procedure described using a TLR3 agonist exhibit greater survival than other patients. One class of patients in which enhanced survival was found were patients having tumors that express a TLR3 protein.

15

Determining whether tumor types express TLR3 can be carried out as described in the examples, e.g. by detecting the presence of one or more TLR3 polypeptides in a biological sample from a cancer patient, generally from a tumor biopsy. The inventors provide herein that several tumor types can express TLR3 proteins and that these types of tumors can be treated with a TLR3 agonist according to the invention.

20

In other specific embodiments, a diagnostic assay is performed on a tumor sample from a patient to determine whether the tumor sample comprises TLR3-expressing cells. Such assays are described herein; for example antibody-based immunohistochemistry assays can be used advantageously. Preferably a tumor biopsy is performed, yielding a biological sample. A determination that said biological sample comprises TLR3 expressing cells indicates that the patient can benefit from the TLR3 agonist administration. The patient is then treated with the TLR3 agonist.

25

Preferably, the step of determining whether cancer cells in said subject express a TLR3 receptor is performed on a tumoral sample derived from a patient. For example, the sample can be a biopsy of the patient's tumor, a cell or tissue culture, etc. Such sample can be obtained by conventional methods. In a particular embodiment, the sample is obtained by non-invasive methods and/or from tissue collections.

30

Therefore, in one embodiment of the methods and uses according to the present invention, the step of determining whether cancer cells in said subject express a TLR3 receptor comprises providing a tumoral sample from the patient and detecting the expression of a TLR3. The expression of a TLR3 may be detected at the nucleic acid level or at the polypeptide level. It will be appreciated that detecting TLR3 can be carried by detecting any known TLR3 allelic variant, including but not limited to allelic variations in the cytoplasmic region of TLR3 gene and in the immediate 5' sequence of the TLR3 gene (see Pirieli et al. (2005) Tissue Antigens 66(2): 125, the disclosure of

35

which is incorporated herein by reference), for example a variant at the C/T polymorphism at position 2593, the C/A polymorphism at position 2642 and the A/G polymorphism at position 2690 in the TLR3 gene.

Various techniques known in the art may be used to detect or quantify TLR3 at the nucleotide level, including sequencing, hybridisation, and amplification. Suitable methods include Southern blot (for DNAs), Northern blot (for RNAs), and fluorescent in situ hybridization (FISH). The detection or quantification of TLR3 at the protein level may be achieved by well-known methods including, but not limited to, gel migration, ELISA, radio-immunoassays (RIA) immuno-enzymatic assays (IEMA), Western Blot or another method for detecting a bound ligand. Protein detection methods require the use of a ligand specific for the TLR3 protein, preferably a specific antibody or a double-stranded RNA molecule. The binding of the antibody or dsRNA molecule may be directly detected if the ligand is labelled with, for example, a fluorescent dye, a radioisotope or another detectable moiety. Alternatively, the binding of the ligand to TLR3 may be indirectly detected through the use of a labelled second reagent that binds to the ligand. Such reagents include labelled anti-immunoglobulin antibodies and enzyme-linked antiimmunoglobulin antibodies that bind to the anti-TLR3 antibody, and labelled nucleic acids that can bind to one or both strands of a dsRNA molecule.

Within the context of this invention, an antibody designates a polyclonal antibody, a monoclonal antibody, as well as fragments or derivatives thereof having substantially the same antigen specificity. Fragments include Fab, Fab'2, CDR regions, etc. Derivatives include single-chain antibodies, humanized antibodies, poly-functional antibodies, etc. TLR3-specific antibodies suitable for use in the present invention are commercially available, such as TLR3 monoclonal antibodies, Ref 12-9039, eBioscience, San Diego, CAUSA; polyclonal antiTLR3, Ref abl3555, Abeam, UK; Clone 40C1285, Imgenex Corp, Biocarta US, San Diego, CA; and Clone TLR3.7, catalog no. HM2096, HyCuit biotechnology B.V., The Netherlands.

The antibody may be an agonistic antibody (as described above), an antagonistic antibody or any other antibody that binds to TLR3. Methods for making any of such antibodies are well-known in the art, as well as described in detail above for agonistic antibodies (with the exception of the screening step which for a diagnostic antibody, would be based on affinity for TLR3 rather than activation of that receptor).

In a specific embodiment, the method comprises contacting a sample from the subject with an antibody specific for the TLR3 protein, and determining the presence of an immune complex. In an alternative embodiment, the expression of a TLR3 receptor in said cancer cell is determined using a TLR3-specific primer or probe. Such primer or probes are designed to specifically hybridise with a TLR3 gene, under suitable hybridisation conditions, thereby allowing detection of a gene or RNA coding for TLR3. A particular embodiment comprises contacting a tumor sample from the patient with a TLR3-specific primer or probe, and determining the existence of a hybrid or

amplification product. The presence (or amount) of TLR3 mRNA in a sample can provide an indication as to the expression of said receptor. Such determination may be accomplished by various techniques known in the art, including through RT-PCR. To that purpose, total RNA is isolated from cancer cells using commercially available kits, such as the RNeasy Mini kit (Qiagen, Valencia, CA). DNase I-treated total RNA (3 µg) is reverse-transcribed by using random primers with RNaseH-free reverse transcriptase (Invitrogen, San Diego, CA). TLR3 can be amplified using specific primers described below. TLR3 5'-CTCAGAAGATTACCAGCCGCC-3¹ (SEQ ID NO 3) 5'-CCATTATGAGACAGATCTAATG-3¹ (SEQ ID No 4)) (see US2003/0165479, the disclosure of which is incorporated herein by reference).

10 Prior to determining expression of TLR3, the sample may be treated to improve availability of TLR3 nucleic acids or polypeptides. Such treatment may include, for instance, a lysis of the cells or tissue (e.g., mechanical, enzymatic or physical).

The invention also relates to a diagnostic kit comprising products and reagents for detecting in a tumoral sample from a subject the expression of a TLR3 gene or polypeptide. Said diagnostic kit
15 according to the present invention comprises any primer, any pair of primers, any nucleic acid probe and/or any ligand, preferably antibody, described in the present invention. Said diagnostic kit according to the present invention can further comprise reagents and/or protocols for performing a hybridization, amplification or antigen-antibody immune reaction.

In addition to the discovery that breast cancer patients treated with TLR3 agonist demonstrate
20 increased survival, it was found that a class of patients in which enhanced survival was found when treated in accordance with the procedure described using a TLR3 agonist were patients having metastatic or recurrent, or aggressive breast cancers. Another class of patients in which enhanced survival was found when treated in accordance with the procedure described using a TLR3 agonist were patients having lymph node positive breast cancers.

25 It is provided that cells from breast cancer samples express TLR3. However, it is also envisaged that samples from subjects having other tumor types will contain cells expressing TLR3, and that these subjects can be treated with TLR3 agonists; it will be appreciated that the skilled person may determine which tumors are suitable for treatment using available methods, including the methods described herein.

30

Upregulation of TLR3 Expression in Tumor Cells

Because the efficacy of the methods of treatment described herein depends upon the expression of TLR3 by the cells of the cancer to be treated, it may be advantageous to stimulate TLR3 expression prior to administering a TLR3 agonist. Thus, according to one embodiment, the invention provides
35 a method of treating a subject suffering from cancer comprising the steps of a) administering to said patient an agent that causes increased expression of TLR3 in a cancer cell; and b) administering to said patient a TLR3 agonist.

Agents that cause upregulation of TLR3 expression are known in that art and include, but are not limited to, inflammatory cytokines such as type I interferons, respiratory syncytial virus and attenuated form thereof, and double-stranded RNA molecules. In addition, TLR3 expression can be upregulated by gene therapy techniques that result in in vivo transformation of cancer cells with an expression vector that encodes TLR3. Preferably, such techniques make use of delivery systems, such as viral vector systems or liposome-based systems that are modified to target cancer cells specifically. Targeting is typically achieved by modifying the outer surface of the delivery system to include a ligand or an antibody specific for the cancer cell surface.

The ability of an agent to cause upregulation of TLR3 expression, particularly in the cells of the cancer to be treated by a TLR3 agonist, may be assayed prior to administration of said agent to a subject. Such an assay is preferably carried out in vitro using a culture of cells derived from the cancer to be treated. Such a culture may be an established cell culture line or a culture of cells derived from a sample of the cancer obtained from the subject.

According to a related embodiment, the invention provides a composition of matter for treating a subject having a cancer comprising: an agent that causes upregulation of TLR3 expression; and a TLR3 agonist, wherein said agent and said agonist are present as separate dosage forms, but are associated with one another. The term "associated with one another" as used herein means that the separate dosage forms are packaged together or otherwise attached to one another such that it is readily apparent that the separate dosage forms are intended to be sold and administered as part of the same regimen. The agent and the TLR3 agonist are preferably packaged together in a blister pack or other multi-chamber package, or as connected, separately sealed containers (such as foil pouches or the like) that can be separated by the user (e.g., by tearing on score lines between the two containers).

According to another related embodiment, the invention provides a kit for determining if an agent is capable of upregulating TLR3 expression in a cancer cell, wherein the kit comprises a cancer cell or cancer cell line which can express TLR3; a TLR3-specific ligand; and a reagent for detecting the binding of said TLR3-specific ligand to a TLR3 expressed by said cancer cell.

Cytotoxic and Tumoricidal TLR3 Ligand Complexes

In one embodiment, the invention provides a composition comprising a TLR3 ligand in association with a cytotoxic agent. The term "cytotoxic agent" as used herein is a molecule that is capable of killing or inhibiting the proliferation of a cancer cell. Such complexes take advantage of the fact that TLR3 is expressed endosomally by dendritic cells, but is present on the surface of cancer cells. Thus, the TLR3 ligand portion of the complex targets the cytotoxic agent specifically to cancer cells with minimal affect on cells of the immune system. The term "in association with," as used herein, means that the two agents are either bound to each other through a covalent and/or non-covalent bond; conjugated to one another (e.g., tethered or otherwise connected to one another

directly or through a linking moiety); or are separate from one another, but present in the same carrier, such as a liposome or other carrier.

The TLR3 ligand component of such compositions may be selected from any TLR3 agonist that directly binds TLR3, any TLR antagonist that directly binds TLR3 or any other molecule that
5 directly binds TLR3. Antibodies to TLR3, double-stranded RNA molecule and small molecule ligands of TLR3 are all potential TLR3 ligands that can be used in such compositions. Non-agonistic antibodies to TLR3 are the preferred ligand component and, as described herein, are commercially available.

The cytotoxic agent in such compositions may be selected from any agent known or being
10 developed for the treatment of cancer, including general antiproliferative agents, chemotherapy agents, tumor-specific antiproliferative agents, cytotoxic agents, cytotoxic inhibitory agents, etc. Examples include, but are not limited to, radioisotopes, toxic proteins, toxic small molecules, such as drugs, toxins, immunomodulators, hormones, hormone antagonists, enzymes, oligonucleotides, enzyme inhibitors, therapeutic radionuclides, angiogenesis inhibitors, chemotherapeutic drugs,
15 vinca alkaloids, anthracyclines, epipodophyllotoxins, taxanes, antimetabolites, alkylating agents, antibiotics, COX-2 inhibitors, SN-38, antimitotics, antiangiogenic and apoptotic agents, particularly doxorubicin, methotrexate, taxol, CPT-11, camptothecins, nitrogen mustards, gemcitabine, alkyl sulfonates, nitrosoureas, triazines, folic acid analogs, pyrimidine analogs, purine analogs, platinum coordination complexes, Pseudomonas exotoxin, ricin, abrin, 5-fluorouridine,
20 ribonuclease (RNase), DNase I, Staphylococcal enterotoxin-A, pokeweed antiviral protein, gelonin, diphtheria toxin, Pseudomonas exotoxin, Pseudomonas endotoxin and others (see, e.g., Remington's Pharmaceutical Sciences, 19th Ed. (Mack Publishing Co. 1995); Goodman and Gilman's The Pharmacological Basis of Therapeutics (McGraw Hill, 2001); Pastan et al. (1986) Cell 47:641; Goldenberg (1994) Cancer Journal for Clinicians 44:43; U.S. Pat. No. 6,077,499;
25 <http://oncologyknowledgebase.com/oksite/TargetedTherapeutics/TTOExhibitM.pdf>, and <http://oncologyknowledgebase.com/oksite/TargetedTherapeutics/TTOExhibitS.pdf>, the entire disclosures of which are herein incorporated by reference). It will be appreciated that a toxin can be of animal, plant, fungal, or microbial origin, or can be created de novo by chemical synthesis. If required, the toxins or other compounds can be linked to the TLR3 ligand directly or indirectly,
30 using any of a large number of available methods depending upon the nature of the TLR3 ligand. For example, an agent can be attached at the hinge region of a TLR3 antibody component via disulfide bond formation, using cross-linkers such as N-succinyl 3-(2-pyridyldithio)propionate (SPDP), or via a carbohydrate moiety in the Fc region of the antibody (see, e.g., Yu et al. (1994) Int. J. Cancer 56: 244; Wong, "Chemistry of Protein Conjugation and Cross-linking," (CRC Press
35 1991); Upeslakis et al., "Modification of Antibodies by Chemical Methods," in Monoclonal Antibodies: Principles And Applications, Birch et al. (eds.), pages 187-230 (Wiley-Liss, Inc. 1995); Price, "Production and Characterization of Synthetic Peptide-Derived Antibodies," in Monoclonal

Antibodies: Production, Engineering And Clinical Application, Ritter et al. (eds.), pages 60-84 (Cambridge University Press 1995); Cattel et al. (1989) Chemistry Today 7:51-58; Delprino et al. (1993) J. Pharm. Sci 82:699-704; Arpicco et al. (1997) Bioconjugate Chemistry 8:3; Reisfeld et al. (1989) Antibody, Immunoconj. Radiopharm. 2:217; the entire disclosures of each of which are
5 herein incorporated by reference).

A toxin or other compound can be conjugated to a dsRNA molecule using a wide variety of well-known techniques. Exemplary U.S. patents that describe the preparation of oligonucleotide conjugates include, for example, U.S. Pat. Nos. 4,828,979; 4,948,882; 5,218,105; 5,525,465; 5,541,313; 5,545,730; 5,552,538; 5,578,717; 5,580,731; 5,580,731; 5,591,584; 5,109,124;
10 5,118,802; 5,138,045; 5,414,077; 5,486,603; 5,512,439; 5,578,718; 5,508,046; 4,587,044; 4,605,735; 4,667,025; 4,752,779; 4,789,737; 4,824,941; 4,835,263; 4,876,335; 4,904,582; 4,958,013; 5,482,830; 5,112,963; 5,214,136; 5,082,830; 5,112,963; 5,214,136; 5,245,022; 5,254,469; 5,258,506; 5,262,536; 5,272,250; 5,292,873; 5,317,098; 5,371,241, 5,391,723; 5,416,203, 5,451,463; 5,510,475; 5,512,667; 5,514,785; 5,565,552; 5,567,810; 5,574,142;
15 5,585,481; 5,587,371; 5,595,726; 5,597,696; 5,599,923; 5,599,928 and 5,688,941, each of which is incorporated by reference herein in its entirety.

When a TLR3 ligand and a cytotoxic agent are present as separate agents in the same vehicle, it is necessary that the TLR3 ligand be at least partially exposed on the surface of the vehicle so that the vehicle will be targeted to the TLR3-expressing tumor cell. Conjugation of antibodies to liposomes
20 to make a delivery vehicle ("immunoliposomes") is well known in the art, e.g., described by Torchilin et al., 1992, FASEB J. 6:2716-2719; Bondas et al., 1999, International J. of Pharmaceutics 181:79-93; Tana et al., 1998, Japanese J. of Cancer Res. 89:1201-1211; or Koning et al., 1999, Biochimica et Biophysica Acta 1420:153-167. Conjugation of other TLR3 ligands to liposomes may be carried out in a similar manner.

25

TLR3 Agonist Formulations and Delivery Systems

The methods of treatment according to this invention comprise the step of administering to a subject a TLR3 agonist. Preferably, the TLR3 agonist is formulated into a pharmaceutically acceptable composition. The pharmaceutically acceptable compositions useful in the invention
30 additionally comprise a pharmaceutically acceptable carriers, adjuvants and vehicles. Pharmaceutically acceptable carriers, adjuvants and vehicles that may be used in the pharmaceutical compositions useful in this invention include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride
35 mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances,

polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol and wool fat.

The pharmaceutical compositions of the invention include those suitable for oral, rectal, nasal, topical (including buccal and sublingual), vaginal or parenteral (including subcutaneous,

5 intramuscular, intravenous and intradermal) administration. In certain embodiments, the compound of the formulae herein is administered transdermally (e.g., using a transdermal patch or iontophoretic techniques). Other formulations may conveniently be presented in unit dosage form, e.g., tablets and sustained release capsules, and in liposomes, and may be prepared by any methods well known in the art of pharmacy. See, for example, Remington's Pharmaceutical Sciences, Mack
10 Publishing Company, Philadelphia, PA (17th ed. 1985).

Such preparative methods include the step of bringing into association with the molecule to be administered ingredients such as the carrier that constitutes one or more accessory ingredients. In general, the compositions are prepared by uniformly and intimately bringing into association the active ingredients with liquid carriers, liposomes or finely divided solid carriers or both, and then if
15 necessary shaping the product.

In certain preferred embodiments, the compound is administered orally. Compositions of the present invention suitable for oral administration may be presented as discrete units such as capsules, sachets or tablets each containing a predetermined amount of the active ingredient; as a powder or granules; as a solution or a suspension in an aqueous liquid or a non-aqueous liquid; or
20 as an oil-in-water liquid emulsion or a water-in-oil liquid emulsion, or packed in liposomes and as a bolus, etc. Soft gelatin capsules can be useful for containing such suspensions, which may beneficially increase the rate of compound absorption.

A tablet may be made by compression or molding, optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing in a suitable machine the active
25 ingredient in a free-flowing form such as a powder or granules, optionally mixed with a binder, lubricant, inert diluent, preservative, surface-active or dispersing agent. Molded tablets may be made by molding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent. The tablets optionally may be coated or scored and may be formulated so as to provide slow or controlled release of the active ingredient therein. Methods of formulating such
30 slow or controlled release compositions of pharmaceutically active ingredients, such as those herein and other compounds known in the art, are known in the art and described in several issued US Patents, some of which include, but are not limited to, US Patent Nos. 4,369,172; and 4,842,866, and references cited therein. Coatings can be used for delivery of compounds to the intestine (see, e.g., U.S. Patent Nos. 6,638,534, 5,217,720, and 6,569,457, 6,461,631, 6,528,080,
35 6,800,663, and references cited therein).

In the case of tablets for oral use, carriers that are commonly used include lactose and corn starch. Lubricating agents, such as magnesium stearate, are also typically added. For oral administration

in a capsule form, useful diluents include lactose and dried cornstarch. When aqueous suspensions are administered orally, the active ingredient is combined with emulsifying and suspending agents. If desired, certain sweetening and/or flavoring and/or coloring agents may be added. Surfactants such as sodium lauryl sulfate may be useful to enhance dissolution and absorption.

- 5 Compositions suitable for oral administration include lozenges comprising the ingredients in a flavored basis, usually sucrose and acacia or tragacanth; and pastilles comprising the active ingredient in an inert basis such as gelatin and glycerin, or sucrose and acacia.
- Compositions suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render
- 10 the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampules and vials, and may be stored in a freeze dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, for example water for injections, immediately prior to use. Extemporaneous
- 15 injection solutions and suspensions may be prepared from sterile powders, granules and tablets. Such injection solutions may be in the form, for example, of a sterile injectable aqueous or oleaginous suspension. This suspension may be formulated according to techniques known in the art using suitable dispersing or wetting agents (such as, for example, Tween 80) and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in
- 20 a non-toxic parenterally-acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are mannitol, water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose, any bland fixed oil may be employed including synthetic mono- or diglycerides. Fatty acids, such as oleic acid and its
- 25 glyceride derivatives are useful in the preparation of injectables, as are natural pharmaceutically-acceptable oils, such as olive oil or castor oil, especially in their polyoxyethylated versions. These oil solutions or suspensions may also contain a long-chain alcohol diluent or dispersant such as Ph. HeIv or a similar alcohol.
- The pharmaceutical compositions of this invention may be administered in the form of
- 30 suppositories for rectal or vaginal administration. These compositions can be prepared by mixing a compound of this invention with a suitable non-irritating excipient which is solid at room temperature but liquid at the rectal temperature and therefore will melt in the rectum to release the active components. Such materials include, but are not limited to, cocoa butter, beeswax and polyethylene glycols.
- 35 Topical administration of the pharmaceutical compositions of this invention is especially useful when the desired treatment involves areas or organs readily accessible by topical application. For application topically to the skin, the pharmaceutical composition will be formulated with a suitable

ointment containing the active components suspended or dissolved in a carrier. Carriers for topical administration of the compounds of this invention include, but are not limited to, mineral oil, liquid petroleum, white petroleum, propylene glycol, polyoxyethylene polyoxypropylene compound, emulsifying wax and water. Alternatively, the pharmaceutical composition can be formulated with

5 a suitable lotion or cream containing the active compound suspended or dissolved in a carrier. Suitable carriers include, but are not limited to, mineral oil, sorbitan monostearate, polysorbate 60, cetyl esters wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water. The pharmaceutical compositions of this invention may also be topically applied to the lower intestinal tract by rectal suppository formulation or in a suitable enema formulation. Topically-transdermal patches and

10 iontophoretic administration are also included in this invention. The pharmaceutical compositions of this invention may be administered by nasal aerosol or inhalation. Such compositions are prepared according to techniques well-known in the art of pharmaceutical formulation and may be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons,

15 and/or other solubilizing or dispersing agents known in the art. Aerosol formulations that may be utilized in the methods of this invention also include those described in United States Patent 6,811,767, the disclosure of which is herein incorporated by reference.

For dsRNA TLR3 agonists, particularly those dsRNA TLR3 agonists that are also siRNA molecules, effective delivery requires that at least a portion of the administered TLR3 agonist enter

20 into the tumor cell. Naked siRNA is known to slowly enter cells by a mechanism that has not yet been elucidated. However, naked siRNA is not very stable in serum and therefore it is preferred that the siRNA compositions useful in this invention be formulated in a delivery vehicle. Preferably the delivery vehicle for a dsRNA TLR3 agonist that is also a siRNA is a liposome. Liposomes fall into two broad classes. Cationic liposomes are positively charged liposomes which

25 interact with the negatively charged dsRNA molecules to form a stable complex. The positively charged dsRNA/liposome complex binds to the negatively charged cell surface and is internalized in an endosome. Due to the acidic pH within the endosome, the liposomes are ruptured, releasing their contents into the cell cytoplasm (Wang et al., Biochem. Biophys. Res. Commun., 1987, 147, 980-985).

30 Liposomes which are pH-sensitive or negatively-charged, entrap dsRNA rather than complex with it. Since both the dsRNA and the lipid are similarly charged, repulsion rather than complex formation occurs. The dsRNA is thus entrapped in the aqueous interior of these liposomes. pH-sensitive liposomes have been used, for example, to deliver dsRNA encoding the thymidine kinase gene to cell monolayers in culture (Zhou et al., Journal of Controlled Release, 1992, 19, 269-274).

35 One major type of liposomal composition includes phospholipids other than naturally-derived phosphatidylcholine. Neutral liposome compositions, for example, can be formed from dimyristoyl phosphatidylcholine (DMPC) or dipalmitoyl phosphatidylcholine (DPPC). Anionic liposome

compositions generally are formed from dimyristoyl phosphatidylglycerol, while anionic fusogenic liposomes are formed primarily from dioleoyl phosphatidylethanolamine (DOPE). Another type of liposomal composition is formed from phosphatidylcholine (PC) such as, for example, soybean PC, and egg PC. Another type is formed from mixtures of phospholipid and/or phosphatidylcholine
5 and/or cholesterol.

Liposomes that include nucleic acids have been described, for example, in Thierry et al., WO 96/40062 (methods for encapsulating high molecular weight nucleic acids in liposomes); Tagawa et al., U.S. Pat. No. 5,264,221 (protein-bonded liposomes containing RNA); Rahman et al., U.S. Pat. No. 5,665,710 (methods of encapsulating oligodeoxynucleotides in liposomes); Love et al., WO
10 97/04787 (liposomes that include antisense oligonucleotides).

Another type of liposome, transfersomes are highly deformable lipid aggregates which are attractive for drug delivery vehicles. (Cevc et al., 1998, Biochim Biophys Acta. 1368(2): 201-15.) Transfersomes may be described as lipid droplets which are so highly deformable that they can penetrate through pores which are smaller than the droplet. Transfersomes are adaptable to the
15 environment in which they are used, for example, they are shape adaptive, self-repairing, frequently reach their targets without fragmenting, and often self-loading. Transfersomes can be made, for example, by adding surface edge-activators, usually surfactants, to a standard liposomal composition.

TLR3 agonists that are shRNA also require entry into the cell because the agent that is
20 administered to the subject is a DNA expression vector that encodes the shRNA. That DNA molecule must enter the cell in order to be transcribed. It is to be understood that the term "TLR3 agonist" includes both a DNA molecule that codes upon transcription for an shRNA that is a dsRNA TLR3 agonist and the transcribed shRNA. The liposomal delivery systems described above are one way in which the DNA encoding a shRNA TLR3 agonist can enter the cell.

25 Alternatively, the DNA encoding a shRNA TLR3 agonist can be prepared in a viral vector system that has the capability of entering into cells. These are well-known in the art and include Madzak et al., J. Gen. Virol., 73: 1533-36 (1992) (papovavirus SV40); Berkner et al., Curr. Top. Microbiol. Immunol., 158: 39-61 (1992) (adenovirus); Moss et al., Curr. Top. Microbiol. Immunol., 158: 25-38 (1992) (vaccinia virus); Muzyczka, Curr. Top. Microbiol. Immunol., 158: 97-123 (1992)
30 (adeno-associated virus); Margulskes, Curr. Top. Microbiol. Immunol., 158: 67-93 (1992) (herpes simplex virus (HSV) and Epstein-Barr virus (EBV)); Miller, Curr. Top. Microbiol. Immunol., 158: 1-24 (1992) (retrovirus); Brandyopadhyay et al., Mol. Cell. Biol., 4: 749-754 (1984) (retrovirus); Miller et al., Nature, 357: 455-450 (1992) (retrovirus); Anderson, Science, 256: 808-813 (1992) (retrovirus); C. Hofmann et al., Proc. Natl. Acad. Sci. USA, 1995; 92, pp. 10099-10103
35 (baculovirus).

The liposomal and virus-based delivery systems discussed above may include a targeting molecule to direct delivery of the TLR3 agonist to the cancer. Such targeting molecules include ligands

and/or antibodies (or binding fragments thereof) that recognize a molecule on the tumor cell surface. One particular tumor cell surface antigen that can be targeted by such delivery systems is TLR3 itself. Preferably, the ligand binds to a site on TLR3 that is separate from and does not interfere with the TLR3 agonist binding site.

5 In certain embodiments of the invention it is desirable that a dsRNA TLR3 agonist not enter into a cell. Without being bound by theory, the inventors believe that in certain instances agonizing TLR3 expressed on the surface of a cancer cell is sufficient to kill the cancer cells without the need to agonize cytoplasmic TLR3. Many other cell types and in particular dendritic cells, express TLR3 only internally, typically in the endosomal membrane. In order to distinguish between
10 different cellular locations for TLR3, the agonist must be prevented from entering into the cells of the subject.

One way to achieve this is to administer the dsRNA TLR3 agonist without a vehicle, such as a liposome, that enhances the ability of the agonist to penetrate cells. The ability of naked dsRNA to enter cells appears to be relatively limited. The mechanism by which naked dsRNA enters cells is
15 unknown, but it is a relatively inefficient process and seems to preferentially occur in tissues such as adipose, liver and parts of the kidney. Thus, administering a dsRNA TLR3 agonist that is not formulated into a liposome or other vehicle that enhances cell penetration will be sufficient to prevent agonism of cytoplasmic TLR3 in many instances.

The treatment of cancer typically is performed in conjunction with a surgical procedure to remove
20 most of all of the cancer from the subject. One advantage of such a surgical procedure is the ability to either implant a device that releases a therapeutic agent directly at the site of the cancer or to administer a therapeutic agent locally to the cancer. Various techniques can be used for providing the subject compositions at the site of interest, such as injection, use of catheters, trocars, projectiles, pluronic gel, stents, sustained drug release polymers or other device which provides for
25 internal access.

Thus, according to another embodiment, the invention provides a method of impregnating or filling an implantable drug release device comprising the step of contacting said drug release device with a TLR3 agonist or composition comprising a TLR3 agonist. Implantable drug release devices include, but are not limited to, biodegradable polymer capsules or bullets, non-degradable,
30 diffusible polymer capsules and biodegradable polymer wafers.

According to another embodiment, the invention provides an implantable drug release device impregnated with or containing a TLR3 agonist or a composition comprising a TLR3 agonist, such that said TLR3 agonist is released from said device and is therapeutically active.

The invention also provides a method of treating a patient suffering from cancer comprising the
35 steps of: surgically removing at least a portion of said cancer; and implanting a drug release device impregnated with or comprising a TLR3 agonist at the site of said cancer. This method optionally includes the step of determining if the cells of said cancer express TLR3 prior to the step of

implanting said drug release device.

TLR3 Agonist Combination Compositions

Treatment with a TLR3 agonist as described herein can optionally advantageously be combined with one or more other therapeutic agents useful in the treatment of cancer. Thus, the TLR3 agonists described herein can be used conjointly, or in combination with, another therapeutic agent useful in the treatment of cancer. The TLR3 agonist compositions described above may thus additionally include other therapeutic agents useful in the treatment of cancer. The other therapeutic agent may be in the same or preferably in a separate container. Such agents include other TLR3 agonists of different molecular composition (i.e., either a different type of molecule or the same type of molecule with a different amino acid or nucleotide sequence or a different chemical formula); cytotoxins, including but not limited to those recited above for use in cytotoxic and tumoricidal TLR3 ligand complexes; cytotoxic and tumoricidal TLR3 ligand complexes; agents that target a tumor antigen or a tumor proliferative protein, such as those disclosed above as targets for siRNA; chemotherapy agents including, but not limited to, cisplatin (CDDP), carboplatin, oxaliplatin, procarbazine, mechlorethamine, cyclophosphamide, camptothecin, ifosfamide, melphalan, chlorambucil, busulfan, nitrosurea, dactinomycin, daunorubicin, doxorubicin, bleomycin, plicomycin, mitomycin, etoposide (VP16), tamoxifen, raloxifene, estrogen receptor binding agents, taxol, gemcitabine, navelbine, farnesyl-protein transferase inhibitors, transplatin, 5-fluorouracil, vincristin, vinblastin and methotrexate, or any analog or derivative variant of the foregoing; therapeutic agents and combination of therapeutic agents for treatment of specific cancers, such as *for breast cancer*, doxorubicin, epirubicin, the combination of doxorubicin and cyclophosphamide (AC), the combination of cyclophosphamide, doxorubicin and 5-fluorouracil (CAF), the combination of cyclophosphamide, epirubicin and 5-fluorouracil (CEF), HerceptinTM, tamoxifen, the combination of tamoxifen and a cytotoxin, taxanes including docetaxel and Paclitaxel, the combination of a taxane plus doxorubicin and cyclophosphamide *for colon cancer*, the combination of 5-FU and leucovorin, the combination of 5FU and levamisole, irinotecan (CPT- 11) or the combination of irinotecan, 5 -FU and leucovorin (IFL) or oxaliplatin; *for prostate cancer*, a radioisotope (i.e., palladium, strontium-89 and Indium), leuprolide or other LHR agonists, nonsteroidal antiandrogens (flutamide, nilutamide, and bicalutamide), steroidal antiandrogens (cyproterone acetate), the combination of leuprolide and flutamide, estrogens such as DES, chlorotrianisene, ethinyl estradiol, conjugated estrogens U.S.P., DES-diphosphate, second-line hormonal therapies such as aminoglutethimide, hydrocortisone, flutamide withdrawal, progesterone, and ketoconazole, low-dose prednisone, or other chemotherapy agents or combination of agent reported to produce subjective improvement in symptoms and reduction in PSA level including docetaxel, paclitaxel, estramustine/docetaxel, estramustine/etoposide, estramustine/vinblastine, and estramustine/Paclitaxel; *for melanoma*: dacarbazine (DTIC),

nitrosoureas such as carmustine (BCNU) and lomustine (CCNU), agents with modest single agent activity including vinca alkaloids, platinum compounds, and taxanes, the Dartmouth regimen (cisplatin, BCNU, and DTIC), interferon alpha (IFN-A), and interleukin-2 (*JL-2*); *for ovarian cancer*. Paclitaxel, docetaxel, cisplatin, oxaliplatin, hexamethylmelamine, tamoxifen, ifosfamide, the combination of paclitaxel (Taxol) or docetaxel (Taxotere) and cisplatin or carboplatin, the combination of cyclophosphamide and cisplatin, the combination of cyclophosphamide and carboplatin, the combination of 5-fluorouracil (5FU) and leucovorin, etoposide, liposomal doxorubicin, gerucitabine or topotecan/*lung cancer*, cisplatin, vincristine, vinblastine, mitomycin, doxorubicin, and etoposide, alone or in combination, the combination of cyclophosphamide, doxorubicin, vincristine/etoposide, and cisplatin (CAV/EP), the combination of cisplatin and vinorelbine, paclitaxel, docetaxel or gemcitabine, and the combination of carboplatin and paclitaxel.

The methods of using a TLR3 agonist composition described herein may also comprises combination treatment with an anti-angiogenic agent. The TLR3 agonist compositions described above may thus also include an anti-angiogenic agent. New blood vessel formation (angiogenesis) is a fundamental event in the process of tumor growth and metastatic dissemination. The vascular endothelial growth factor (VEGF) pathway is well established as one of the key regulators of this process. The VEGF/VEGF-receptor axis is composed of multiple ligands and receptors with overlapping and distinct ligand-receptor binding specificities, cell-type expression, and function. Activation of the VEGF-receptor pathway triggers a network of signaling processes that promote endothelial cell growth, migration, and survival from pre-existing vasculature. In addition, VEGF mediates vessel permeability, and has been associated with malignant effusions. The VEGF-related gene family comprises six secreted glycoproteins referred to as VEGF-A, VEGF-B, VEGF-C, VEGF-D, VEGF-E, and placenta growth factor (PlGF)-1 and -2. A number of exemplary anti-angiogenic agents acting on the VEGF pathway are known, any of which can be used in accordance with the invention, including small molecule inhibitor, neutralizing antibodies antisense strategies, RNA aptamers and ribozymes against VEGF-related gene family (e.g. the VEGF-A, VEGF-B, VEGF-C, VEGF-D, VEGF-E proteins). Variants of VEGF with antagonistic properties may also be employed, as described in WO 98/16551, specifically incorporated herein by reference. Further exemplary anti-angiogenic agents that are useful in connection with combined therapy are listed in Table D of U.S. Patent No. 6,524,583, the disclosure of which agents and indications are specifically incorporated herein by reference. Particularly preferred anti-angiogenic agents inhibit signaling by a receptor tyrosine kinase including but not limited to VEGFR1, VEGFR-2,3 PDGFR-beta, Flt-3, c-Kit, p38 alpha and FGFR-1. Further anti-angiogenic agent may include agents that inhibit one or more of the various regulators of VEGF expression and production, such as EGFR, HER-2, COX-2, or HIF-1. Another preferred class of agents includes thalidomide or the analogue CC-5013.

- Selected VEGF/VEGFR inhibitors of the monoclonal antibody (mAb) and tyrosine kinase inhibitor (TKI) class include bevacuzimab (Avastin) (mAb, inhibiting VEGF-A, Genentech); IMC-1 121B (mAb, inhibiting VEGFR-2, ImClone Systems); CDP-791 (Pegylated DiFab, VEGFR-2, Celltech); 2C3 (mAb, VEGF-A, Peregrine Pharmaceuticals); PTK-787 (TKI, VEGFR-I, -2, Novartis);
- 5 AEE788 (TKI, VEGFR-2 and EGFR, Novartis); ZD6474 (TKI, VEGFR-I, -2, -3, EGFR AstraZeneca); AZD2171 (TKI, VEGFR-I, -2, AstraZeneca); SU1 1248 (TKI, VEGFR-I, -2, PDGFR Pfizer); AG13925 (TKI, VEGFR-I, -2, Pfizer); AG013736 (TKI, VEGFR-I, -2, Pfizer); CEP-7055 (TKI, VEGFR-I, -2, -3, Cephalon); CP-547,632 (TKI, VEGFR-I, -2, Pfizer); VEGF-trap (Soluble hybrid receptor VEGF-A, PlGF (placenta growth factor) Aventis/Regeneron);
- 10 GW786024 (TKI, VEGFR-I, -2, -3, GlaxoSmithKline); Bay 93-4006 (TKI, VEGFR-I, -2, PDGFR Bayer/Onyx); and AMG706 (TKI, VEGFR-I, -2, -3, Amgen). Most preferred are tyrosine kinase inhibitors that inhibit one or more receptor tyrosine kinases selected from the group consisting of VEGFR1, VEGFR-2,3 PDGFR-beta, Flt-3, c-Kit, p38 alpha and FGFR-I. Preferred examples include SU1 1248 (Pfizer) and Bay 93-4006 (sorefanib, Bayer).
- 15 The methods of using a TLR3 agonist composition described herein may also comprises combination treatment with a pro-apoptotic agent. The TLR3 agonist compositions described above may thus also include a pro-apoptotic agent. A number of proteins useful as targets for modulation by pharmaceutical agents are known in the art, including any of those reviewed in Green and Rroemer, J. (2005) Clin. Investig. 115(10):2610-2617, the disclosure of which is incorporated by
- 20 reference in its entirety. Examples of pro-apoptotic pharmaceutical agents include:
- drugs that induce mitochondrial outer membrane permeabilization such as oblimirsen (Bcl-2 antisense oligonucleotide, Genta), EGCG (small molecule targeting Bcl-2, Burnham Inst/Mayo Clinic), Gossypol (small molecule targeting Bcl-2, Univ. Michigan), LY2181308 (antisense oligonucleotide targeting surviving (Eli Lilly/Isis),
- 25 and arsenic trioxide;
- drugs that regulate p53 activity such as Advexin INGN201 (adenovirus modulating p53, Introgen), SCH58500 (adenovirus modulating p53, Schering-Plough), ONYX-015 (ElB mutated adenovirus modulating p53, Onyx Pharma.);
- drugs that modulate caspases and/or endogenous inhibitors or caspases such as
- 30 AEG35 156 (antisense oligonucleotide targeting XIAP, Aegara/Hybridon);
- drugs that modulate death receptors and/or their ligands, such as TNF-alpha polypeptides, HGS-ETR1 (agonistic mAb targeting TRAIL-R1, Human Genome Sciences), HGS-ETR2 and HGS-TR2J ((two agonistic mAbs targeting TRAIL-R2, Human Genome Sciences); PRO1 764 (soluble TRAIL ligands, Genentech/Amgen);
- 35 and
- drugs targeting poly(ADP-ribose) polymerase (PARP), such as AGO14699 (small molecule, Cancer Res. Tech.);

- drugs targeting the proteasome, such as bortezomib (Velcade) (26S proteasome inhibitor, Millennium Pharma.); and
- kinase inhibitors, such as herceptin (mAb targeting HER2, Roche), centuximab (mAb targeting HER1, Inclone/BMS), gefitinib (Iressa) and erlotinib (Tarceva) (small molecule inhibitors of HER1, AstraZeneca and Genentech/OSI respectively, CCI-779 (small molecule acting on mTOR, Novartis, Bay 43-9006 (small molecule inhibitor of kinases including Raf and VEGFR, and imatinib mesylate (Gleevec, STI-571) (small molecule inhibitor of cKit, PDGFR, Bcr-Abl, Novartis).
- 10 The TLR3 agonist compositions described above may also include other therapeutic agents such as immunomodulatory agents such as tumor necrosis factor, interferon alpha, beta, and gamma, EL-2, IL-12, IL-15, IL-21, CpG-containing single-stranded DNA, agonists of other TLRs, other cytokines and immunosuppression agents; F42K and other cytokine analogs; or MIP-1, MIP-1beta, MCP-1, RANTES, and other chemokines; agents that affect the upregulation of cell surface receptors and
- 15 GAP junctions; cytostatic and differentiation agents; or inhibitors of cell adhesion.
- It will be recognized by those of skill in the art that certain therapeutic agents set forth above fall into two or more of the categories disclosed above. For the purpose of this invention, such therapeutic agents are to be considered members of each of those categories of therapeutics and the characterization of any therapeutic agent as being in a certain specified category does not preclude
- 20 it from also being considered to be within another specified category.
- In yet another embodiment, the invention provides a composition of matter comprising a TLR3 agonist and another therapeutic agent useful in the treatment of cancer in separate dosage forms, but associated with one another. The term "associated with one another" as used herein means that the separate dosage forms are packaged together or otherwise attached to one another such that it is
- 25 readily apparent that the separate dosage forms are intended to be sold and administered as part of the same regimen. The agent and the TLR3 agonist are preferably packaged together in a blister pack or other multi-chamber package, or as connected, separately sealed containers (such as foil pouches or the like) that can be separated by the user (e.g., by tearing on score lines between the two containers).
- 30 In another embodiment, a dsRNA TLR3 agonist may be formulated together in a composition with an endocytosis inhibitor. Dendritic cells express TLR3 on their endosomal surface and are believed to take up dsRNA through endocytosis. Agonism of TLR3 in dendritic cells causes maturation of dendritic cells, production of type I interferons and the secretion of various cytokines. In a tumor environment, certain of those cytokines may cause tumor growth. Thus, in
- 35 certain embodiments it is desirable to prevent a dsRNA TLR3 agonist from entering into dendritic cells. Compositions comprising a dsRNA TLR3 agonist and an agent that blocks endocytosis can be used to treat cancer characterized by a TLR3-expressing cancer cell without stimulating TLR3

in dendritic cells. Examples of an agent that inhibits endocytosis include, but are not limited to, an actin stabilizing agent, such as latrunculin B, and an endosomal pH neutralizing agent, such as chloroquine or hydroxychloroquine.

In a related embodiment, the endocytosis inhibitor may be conjugated to the dsRNA TLR3 agonist.

- 5 The two agents may be either covalently bonded or conjugated directly to one other or attached via a linker or tether moiety.

In yet another related embodiment, a dsKNA TLR3 agonist and an endocytosis inhibitor may be present in separate dosage forms, but associated with one another.

10 **Cancer and Therapeutic Methods**

The present invention encompasses treatment protocols that provide better prophylactic or therapeutic profiles than current single agent therapies or combination therapies for cancer. In particular, the invention encompasses the use of an TLR3 agonist for the manufacture of a medicament for the prevention, management, treatment or amelioration of cancer or one or more
15 symptoms thereof.

Examples of cancers that can be prevented, managed, treated or ameliorated in accordance with the methods invention include, but are not limited to, solid tumors, and particularly cancers such as cancer of the head, neck, eye, mouth, throat, esophagus, chest, bone, lung, colon, rectum, stomach, prostate, breast, ovaries, kidney, liver, pancreas, and brain.

- 20 The invention provides methods for preventing, managing, treating or ameliorating cancer that has the potential to metastasize or has metastasized to an organ or tissue (e.g., bone) or one or more symptoms thereof, said methods comprising administering to a subject in need thereof one or more doses of a prophylactically or therapeutically amount of a TLR3 agonist. The invention also concerns the use of an TLR3 agonist for the manufacture of a medicament for preventing,
25 managing, treating or ameliorating cancer that has the potential to metastasize or has metastasized to an organ or tissue (e.g., bone) or one or more symptoms thereof. Preferably, the TLR3 agonist is a dsRNA compound. Preferably, the TLR3 agonist is administered more than once. Optionally, the TLR3 agonist is administered at an interval of less than one month, less than three weeks, less than two weeks, or less than one week. Optionally, such treatment may be repeated, for example, every
30 1, 2, 3, 4, 5, 6, or 7 days, or every 1, 2, 3, 4, and 5 weeks or every 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12 months.

The present invention provides methods for preventing, managing, treating or ameliorating cancer or one or more symptoms thereof, said methods comprising administering to a subject in need thereof a dosage of a prophylactically or therapeutically effective amount of a TLR3 agonist in
35 combination with the administration of a dosage of a prophylactically or therapeutically effective amount of one or more other agents useful for cancer therapy. The invention also concerns the use of an TLR3 agonist for the manufacture of a medicament for preventing, managing, treating or

ameliorating cancer, wherein said TLR3 agonist is used in combination with one or more agents useful for cancer therapy. Preferably, the agonist is a dsRNA compound. Preferably, the TLR3 agonist is administered more than once. Optionally, the TLR3 agonist is administered at an interval of less than one month, less than three weeks, less than two weeks, or less than one week.

5 Optionally, such treatment may be repeated, for example, every 1, 2, 3, 4, 5, 6, or 7 days, or every 1, 2, 3, 4, and 5 weeks or every 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12 months.

The invention provides methods for preventing, managing, treating or ameliorating cancer that has been refractory to one or more therapeutic agents or therapies, said methods comprising administering to a subject in need thereof one or more doses of a prophylactically or therapeutically
10 amount of a TLR3 agonist. The invention also concerns the use of an TLR3 agonist for the manufacture of a medicament for preventing, managing, treating or ameliorating cancer that has been refractory to one or more therapeutic agents or therapies. Preferably, the TLR3 agonist is a dsRNA compound. Preferably, the doses are administered at an interval of less than one month, less than three weeks, less than two weeks, or less than one week. Optionally, such treatment may
15 be repeated, for example, every 1, 2, 3, 4, 5, 6, or 7 days, or every 1, 2, 3, 4, and 5 weeks or every 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12 months.

The present invention provides methods for preventing, treating, managing or ameliorating cancer or one or more symptoms thereof, said methods comprising administering to a subject in need thereof a TLR3 agonist alone or in combination with one or more other therapies (e.g., one or more
20 other prophylactic or therapeutic agents) useful in the prevention, treatment, management or amelioration of cancer or one or more symptoms thereof. Other therapies useful in combination with the administration of TLR3 agonist also include treatment with any of the agents described above that are useful in TLR3 agonist combination compositions.

In all of the treatment regimens it is preferred that the TLR3 agonist is a dsRNA compound.
25 Treatments involving the administration of a TLR3 agonist in combination with one or more therapies include simultaneous administration and treatment either as separate therapies administered at the same time (such as multiple dosage forms or a physical treatment such as exposure to radiation, while administering the TLR3 agonist), or as a single combination therapy (such as the combination single dosage forms described above containing a TLR3 agonist and
30 another therapeutic agent. A treatment involving the administration of a TLR3 agonist in combination with one or more therapies also includes consecutive or staggered treatments where the TLR3 agonist is administered at one point in time and the other therapy is administered from 1 minute to 1 month before or after that time.

In one embodiment, a TLR3 agonist (preferably, a dsRNA) is administered to a subject using a
35 dosing regimen that maintains the plasma concentration of the agonist at a desirable level. In a specific embodiment, the plasma concentration of the dsRNA is maintained at 10 µg/ml, 15 µg/ml, 20 µg/ml, 25 µg/ml, 30 µg/ml, 35 µg/ml, 40 µg/ml, 45 µg/ml or 50 µg/ml. The plasma

concentration that is desirable in a subject will vary depending on several factors including, but not limited to, the nature of the cancer, the severity of the cancer, and the circulation half-life (stability) and binding affinity of the TLR3 agonist.

5 The dosage amounts and frequencies of administration provided herein are encompassed by the terms therapeutically effective and prophylactically effective. The dosage and frequency further will typically vary according to factors specific for each patient depending on the specific therapeutic or prophylactic agents administered, the severity and type of cancer, the route of administration, as well as age, body weight.

10 Preferably, a therapeutically effective amount of a TLR3 agonist (optionally in combination with another therapeutic agent or therapeutic protocol) reduces the size of a tumor or the spread of a tumor in a subject by at least 5%, preferably at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95% or at least 99% relative to a control such as PBS.

15 As used herein, the terms "non-responsive" and "refractory" describe patients treated with a currently available cancer therapy (e.g., chemotherapy, radiation therapy, surgery, hormonal therapy and/or biological therapy/immunotherapy), which is not clinically adequate to treat or relieve one or more symptoms associated with cancer. Typically, such patients suffer from severe, persistently active disease and require additional therapy to ameliorate the symptoms associated with their cancer. The phrase can also describe patients who respond to therapy yet suffer from side effects, relapse, develop resistance, etc.

20 In various embodiments, "non-responsive/refractory" means that at least some significant portion of the cancer cells are not killed or their cell division arrested. The determination of whether the cancer cells are "non-responsive/refractory" can be made either in vivo or in vitro by any method known in the art for assaying the effectiveness of treatment on cancer cells, using the art-accepted meanings of "refractory" in such a context. In various embodiments, a cancer is "non-responsive/refractory" when the number of cancer cells has not been significantly reduced, or has increased.

Types of cancers

30 In various embodiments, the present invention provides methods for determining treatment regimens for cancer subjects. The methods of the invention can be used to determine treatment regimens of any cancer, or tumor, for example, but not limited to, malignancies and related disorders include but are not limited to the following.

35 Leukemias such as but not limited to, acute leukemia, acute lymphocytic leukemia, acute myelocytic leukemias such as myeloblastic, promyelocytic, myelomonocytic, monocytic, erythroleukemia leukemias and myelodysplastic syndrome, chronic leukemias such as but not limited to, chronic myelocytic (granulocytic) leukemia, chronic lymphocytic leukemia, hairy cell

leukemia; polycythemia vera; lymphomas such as but not limited to Hodgkin's disease, non-Hodgkin's disease; multiple myelomas such as but not limited to smoldering multiple myeloma, nonsecretory myeloma, osteosclerotic myeloma, plasma cell leukemia, solitary plasmacytoma and extramedullary plasmacytoma; Waldenstrom's macroglobulinemia; monoclonal gammopathy of

5 undetermined significance; benign monoclonal gammopathy; heavy chain disease; bone and connective tissue sarcomas such as but not limited to bone sarcoma, osteosarcoma, chondrosarcoma, Ewing's sarcoma, malignant giant cell tumor, fibrosarcoma of bone, chordoma, periosteal sarcoma, soft-tissue sarcomas, angiosarcoma (hemangiosarcoma), fibrosarcoma, Kaposi's sarcoma, leiomyosarcoma, liposarcoma, lymphangiosarcoma, neurilemmoma,

10 rhabdomyosarcoma, synovial sarcoma; brain tumors such as but not limited to, glioma, astrocytoma, brain stem glioma, ependymoma, oligodendroglioma, nonglial tumor, acoustic neurinoma, craniopharyngioma, medulloblastoma, meningioma, pineocytoma, pineoblastoma, primary brain lymphoma; breast cancer including but not limited to adenocarcinoma, lobular (small cell) carcinoma, intraductal carcinoma, medullary breast cancer, mucinous breast cancer, tubular

15 breast cancer, papillary breast cancer, Paget's disease, and inflammatory breast cancer; adrenal cancer such as but not limited to pheochromocytoma and adrenocortical carcinoma; thyroid cancer such as but not limited to papillary or follicular thyroid cancer, medullary thyroid cancer and anaplastic thyroid cancer; pancreatic cancer such as but not limited to, insulinoma, gastrinoma, glucagonoma, vipoma, somatostatin-secreting tumor, and carcinoid or islet cell tumor; pituitary

20 cancers such as but limited to Cushing's disease, prolactin-secreting tumor, acromegaly, and diabetes insipidus; eye cancers such as but not limited to ocular melanoma such as iris melanoma, choroidal melanoma, and ciliary body melanoma, and retinoblastoma; vaginal cancers such as squamous cell carcinoma, adenocarcinoma, and melanoma; vulvar cancer such as squamous cell carcinoma, melanoma, adenocarcinoma, basal cell carcinoma, and sarcoma; cervical cancers such

25 as but not limited to, squamous cell carcinoma, and adenocarcinoma; uterine cancers such as but not limited to endometrial carcinoma and uterine sarcoma; ovarian cancers such as but not limited to, ovarian epithelial carcinoma, borderline tumor, germ cell tumor, and stromal tumor; esophageal cancers such as but not limited to, squamous cancer, adenocarcinoma, adenoid cystic carcinoma, mucoepidermoid carcinoma, adenosquamous carcinoma, sarcoma, melanoma, plasmacytoma,

30 verrucous carcinoma, and oat cell (small cell) carcinoma; stomach cancers such as but not limited to, adenocarcinoma, ulcerating, superficial spreading, diffusely spreading, malignant lymphoma, liposarcoma, fibrosarcoma, and carcinosarcoma; colon cancers; rectal cancers; liver cancers such as but not limited to hepatocellular carcinoma and hepatoblastoma, gallbladder cancers such as adenocarcinoma; cholangiocarcinomas such as but not limited to papillary, nodular, and diffuse;

35 lung cancers such as non-small cell lung cancer, squamous cell carcinoma (epidermoid carcinoma), adenocarcinoma, large-cell carcinoma and small-cell lung cancer; testicular cancers such as but not limited to germinal tumor, seminoma, anaplastic, classic (typical), spermatocytic, nonseminoma,

embryonal carcinoma, teratoma carcinoma, choriocarcinoma (yolk-sac tumor), prostate cancers such as but not limited to, adenocarcinoma, leiomyosarcoma, and rhabdomyosarcoma; penile cancers; oral cancers such as but not limited to squamous cell carcinoma; basal cancers; salivary gland cancers such as but not limited to adenocarcinoma, mucoepidermoid carcinoma, and

5 adenoidcystic carcinoma; pharynx cancers such as but not limited to squamous cell cancer, and venaceous; skin cancers such as but not limited to, basal cell carcinoma, squamous cell carcinoma and melanoma, superficial spreading melanoma, nodular melanoma, lentigo malignant melanoma, acral lentiginous melanoma; kidney cancers such as but not limited to renal cell cancer, adenocarcinoma, hypernephroma, fibrosarcoma, transitional cell cancer (renal pelvis and/ or

10 ureter); Wilkins' tumor; bladder cancers such as but not limited to transitional cell carcinoma, squamous cell cancer, adenocarcinoma, carcinosarcoma. In addition, cancers include myxosarcoma, osteogenic sarcoma, endotheliosarcoma, lymphangioendotheliosarcoma, mesothelioma, synovioma, hemangioblastoma, epithelial carcinoma, cystadenocarcinoma, bronchogenic carcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma

15 and papillary adenocarcinomas (for a review of such disorders, see Fishman et al., 1985, Medicine, 2d Ed., J.B. Lippincott Co., Philadelphia and Murphy et al., 1997, Informed Decisions: The Complete Book of Cancer Diagnosis, Treatment, and Recovery, Viking Penguin, Penguin Books U.S.A., Inc., United States of America).

Accordingly, the methods of the invention are also useful in the treatment of a variety of cancers or ,

20 other abnormal proliferative diseases, including (but not limited to) the following: carcinoma, including that of the bladder, breast, colon, kidney, liver, lung, ovary, pancreas, stomach, cervix, thyroid and skin; including squamous cell carcinoma; hematopoietic tumors of lymphoid lineage, including leukemia, acute lymphocytic leukemia, acute lymphoblastic leukemia, B-cell lymphoma, T-cell lymphoma, Berketts lymphoma; hematopoietic tumors of myeloid lineage, including acute

25 and chronic myelogenous leukemias and promyelocytic leukemia; tumors of mesenchymal origin, including fibrosarcoma and rhabdomyosarcoma; other tumors, including melanoma, seminoma, teratocarcinoma, neuroblastoma and glioma; tumors of the central and peripheral nervous system, including astrocytoma, neuroblastoma, glioma, and schwannomas; tumors of mesenchymal origin, including fibrosarcoma, rhabdomyosarcoma, and osteosarcoma; and other tumors, including

30 melanoma, xenoderma pigmentosum, keratoacanthoma, seminoma, thyroid follicular cancer and teratocarcinoma. In specific embodiments, malignancy or dysproliferative changes (such as metaplasias and dysplasias), or hyperproliferative disorders, are treated in the ovary, bladder, breast, colon, lung, skin, pancreas, or uterus. In other specific embodiments, sarcoma, melanoma, or leukemia is treated.

35 In preferred embodiments, the methods of the invention are used for TLR3 positive solid tumors. Example of tumors include breast, colon, ovarian, lung, brain and prostate cancers and melanoma. In a preferred embodiment, the methods of the invention are directed at treating breast cancer.

Breast Cancer

Types of Breast Cancer. Most breast cancer develops in glandular tissue and is classified as adenocarcinoma. The earliest form of the disease, ductal carcinoma in situ (DCIS), develops solely in the milk ducts. The most common type of breast cancer, invasive ductal carcinoma (IDC),

5 develops from DCIS, spreads through the duct walls, and invades the breast tissue.

The term "pre-malignant lesion" as used herein is defined as a collection of cells in a breast with histopathological characteristics which suggest at least one of the cells has an increased risk of becoming breast cancer. A skilled artisan recognizes that the most important pre-malignant lesions recognized today include unfolded lobules (UL; other names: blunt duct adenosis, columnar

10 alteration of lobules), usual ductal hyperplasia (UDH; other names: proliferative disease without atypia, epitheliosis, papillomatosis, benign proliferative disease), atypical ductal hyperplasia (ADH), atypical lobular hyperplasia (ALH), ductal carcinoma in situ (DCIS), and lobular carcinoma in situ (LCIS). Other lesions which may have pre-malignant potential include intraductal papillomas, sclerosing adenosis, and fibroadenomas (especially atypical fibroadenomas). In a

15 specific embodiment, the collection of cells is a lump, tumor, mass, bump, bulge, swelling, and the like. Other terms in the art which are interchangeable with "pre-malignant lesion" include pre-malignant hyperplasia, pre-malignant neoplasia, and the like.

Invasive lobular carcinoma originates in the milk glands and accounts for 10 -15% of invasive breast cancers. Less common types of breast cancer include the following:

20 Inflammatory (breast tissue is warm and appears red; tends to spread quickly)

Medullary carcinoma (originates in central breast tissue)

Mucinous carcinoma (invasive; usually occurs in postmenopausal women)

Paget's disease of the nipple (originates in the milk ducts and spreads to the skin of the nipples or areola)

25 Phyllodes tumor (tumor with a leaf-like appearance that extends into the ducts; rarely metastasizes)

Tubular carcinoma (small tumor that is often undetectable by palpation)

Rarely, sarcomas (cancer of the connective tissue) and lymphomas (cancer of the lymph tissue) develop in the breasts.

30 *Staging* : The stage of a cancer is determined by the size and location in the body of the primary tumor, and whether it has spread to other areas of the body.

Staging involves using the letters T, N, and M to assess tumors:

size of the primary tumor (T);

degree to which regional lymph nodes (N) are involved. Lymph nodes are small organs

35 located along the channels of the body's lymphatic system which store special cells that fight infection and other diseases); and

absence or presence of distant metastases (M) - cancer that has spread from the original (primary) tumor to distant organs or distant lymph nodes.

Each of these categories is further classified with a number. Thus a T1-N1-M0 cancer would describe a T1 tumor, N1 lymph node involvement, and no metastases.

- 5 Once the T, N, and M components are determined, a "stage" of I, II, III or IV is assigned:

Stage I cancers are small, localized and usually curable.

Stage II and III cancers typically are locally advanced and/or have spread to local lymph nodes.

- 10 Stage IV cancers usually are metastatic (have spread to distant parts of the body) and generally are considered inoperable.

Details of this staging system are further provided in: The International Union against Cancer details the TNM staging system (Tumour/Nodes/Metastasis), (UICC - TNM Classification of malignant tumours. Edited by L.H. Sobin and CH. Wittekind. 5th Edition. New York: Wiley-Liss; 1997)

- 15 The term "sample from a breast" as used herein is defined as a specimen from any part or tissue of a breast. A skilled artisan recognizes that the sample may be obtained by any method, such as biopsy. In another specific embodiment, the sample is from hyperplastic or malignant breast epithelium. In a specific embodiment, the sample is from the epithelium. In another specific embodiment, the sample is from a premalignant lesion.

- 20 In specific embodiments, a patient with breast cancer is administered a prophylactically or therapeutically effective amount of a TLR3 agonist. The patient may or may not have lymph node involvement, may or may not have metastases to distant organs or distant lymph nodes, may or may not have a recurrent cancer, or a cancer that is refractory or non-refractory. The cancer may be a stage I, II, III or IV cancer, most preferably a stage II or III cancer. The cancer may be a T1,

- 25 T2 or T3 cancer. Optionally, the cancer is a T1-3N0-3M0 cancer. Optionally, the patient has received radiation therapy following surgery to remove breast cancer tissue, and in another aspect, the patient has not received radiation therapy following surgery to remove breast cancer tissue.

- Generally, although the therapeutic methods are not limited thereto, patients will have received surgery to remove breast cancer tissue. The TLR3 agonist may also be used as a prophylactic agent before surgery, or more preferably is used in combination with surgery. The TLR3 agonist can advantageously be used soon or immediately after surgery (e.g beginning of TLR3 agonist treatment less than 8, 6, 4, 3, 2, 1 week following surgery), without possible adverse effects on TLR3 agonist efficacy related to immunosuppression that could be expected with immunomodulatory compounds.

- 35 In other specific embodiments, a diagnostic assay is performed on a sample from a breast to determine whether a patient's tumor comprises TLR3-expression cells. Such assays are described herein; for example antibody-based immunohistochemistry assays can be used advantageously.

Preferably a tumor biopsy is performed, yielding a biological sample. A determination that said biological sample comprises TLR3 expressing cells indicates that the patient can benefit from the TLR3 agonist administration. The patient is then treated with the TLR3 agonist.

5 In other specific embodiments, patients with breast cancer are administered a prophylactically or therapeutically effective amount of a TLR3 agonist in combination with a prophylactically or therapeutically effective amount of one or more other therapies useful for breast cancer treatment or management including, but not limited to: doxorubicin, epirubicin, the combination of doxorubicin and cyclophosphamide (AC), the combination of cyclophosphamide, doxorubicin and 5-fluorouracil (CAF), the combination of cyclophosphamide, epirubicin and 5-fluorouracil (CEF),
10 Herceptin™), tamoxifen, or the combination of tamoxifen and cytotoxic chemotherapy. In certain embodiments, patients with metastatic breast cancer are administered a prophylactically or therapeutically effective amount of a TLR3 agonist in combination with the administration of a prophylactically or therapeutically effective amount of taxanes such as docetaxel and paclitaxel. In other embodiments, a patients with node-positive, localized breast cancer are administered a
15 prophylactically or therapeutically effective amount of a TLR3 agonist in combination with the administration of a prophylactically or therapeutically effective amount of taxanes plus standard doxorubicin and cyclophosphamide for adjuvant treatment of node-positive, localized breast cancer.

Treatment of Colon Cancer

20 In specific embodiments, a patient with colon cancer is administered a prophylactically or therapeutically effective amount of a TLR3 agonist. The patient may or may not have metastases to distant organs or distant lymph nodes, may or may not have a recurrent cancer, or a cancer that is refractory or non-refractory. Optionally, the patient has received radiation therapy following surgery to remove cancerous tissue. In another aspect, the patient has not received radiation therapy
25 following surgery to remove cancerous tissue.

Generally, although the therapeutic methods are not limited thereto, patients will have received surgery to remove cancer tissue. The TLR3 agonist may also be used a prophylactic agent before surgery, or more preferably is used in combination with surgery. The TLR3 agonist can advantageously be used soon or immediately after surgery (e.g. beginning of TLR3 agonist
30 treatment less than 8, 6, 4, 3, 2, 1 week following surgery), without possible adverse effects on TLR3 agonist efficacy related to immunosuppression.

In specific embodiments, patients with colon cancer are administered a prophylactically or therapeutically effective amount of a TLR3 agonist in combination with the administration of a prophylactically or therapeutically effective amount of one or more other therapies useful for colon
35 cancer treatment or management including but not limited to: the combination of 5-FU and leucovorin, the combination of 5FU and levamisole, irinotecan (CPT- 11) or the combination of irinotecan, 5 -FU and leucovorin (IFL) or oxaliplatin.

Treatment of Prostate Cancer

In specific embodiments, patients with prostate cancer are administered a prophylactically or therapeutically effective amount of a TLR3 agonist in combination with the administration of a prophylactically or therapeutically effective amount of one or more other therapies useful for prostate cancer treatment or management including but not limited to: external-beam radiation therapy, interstitial implantation of radioisotopes (i.e., palladium, and Indium), leuprolide or other LHR agonists, nonsteroidal antiandrogens (flutamide, nilutamide, and bicalutamide), steroidal antiandrogens (cyproterone acetate), the combination of leuprolide and flutamide, estrogens such as DES, chlorotrianisene, ethinyl estradiol, conjugated estrogens U.S.P., DES-diphosphate, radioisotopes, such as strontium-89, the combination of external-beam radiation therapy and strontium-89, second-line hormonal therapies such as aminoglutethimide, hydrocortisone, flutamide withdrawal, progesterone, and ketoconazole, low-dose prednisone, or other chemotherapy regimens reported to produce subjective improvement in symptoms and reduction in PSA level including docetaxel, paclitaxel, estramustine/docetaxel, estramustine/etoposide, estramustine/vinblastine, and estramustine/paclitaxel.

Treatment of Melanoma

In specific embodiments, patients with melanoma are administered a prophylactically or therapeutically effective amount of a TLR3 agonist in combination with the administration of a prophylactically or therapeutically effective amount of one or more other therapies useful for melanoma cancer treatment or management including but not limited to: dacarbazine (DTIC), nitrosoureas such as carmustine (BCNU) and lomustine (CCNU), agents with modest single agent activity including vinca alkaloids, platinum compounds, and taxanes, the Dartmouth regimen (cisplatin, BCNU, and DTIC), interferon alpha (IFN-A), and interleukin-2 (IL-2).

Treatment of Ovarian Cancer

In specific embodiments, patients with ovarian cancer are administered a prophylactically or therapeutically effective amount of a TLR3 agonist in combination with a prophylactically or therapeutically effective amount of one or more other therapies useful for ovarian cancer treatment or management including, but not limited to: intraperitoneal radiation therapy, total abdominal and pelvic radiation therapy, cisplatin, oxaliplatin, the combination of paclitaxel (Taxol) or docetaxel (Taxotere) and cisplatin or carboplatin, the combination of cyclophosphamide and cisplatin, the combination of cyclophosphamide and carboplatin, the combination of 5-fluorouracil (5FU) and leucovorin, etoposide, liposomal doxorubicin, gerucitabine or topotecan. In a particular embodiment, patients with ovarian cancer that is platinum-refractory are administered a prophylactically or therapeutically effective amount of a TLR3 agonist in combination with the administration of a prophylactically or therapeutically effective amount of Taxol. The invention encompasses the treatment of patients with refractory ovarian cancer including administration of ifosfamide in patients with disease that is platinum-refractory, hexamethylmelamine (HAM) as

salvage chemotherapy after failure of cisplatin-based combination regimens, and tamoxifen in patients with detectable levels of cytoplasmic estrogen receptor on their tumors.

Treatment of Lung Cancers

hi specific embodiments, patients with small lung cell cancer are administered a prophylactically or therapeutically effective amount of a TLR3 agonist in combination with the administration of a prophylactically or therapeutically effective amount of one or more other therapies useful for lung cancer treatment or management including but not limited to: thoracic radiation therapy, cisplatin, vincristine, doxorubicin, and etoposide, alone or in combination, the combination of cyclophosphamide, doxorubicin, vincristine/etoposide, and cisplatin (CAVTEP), local palliation with endobronchial laser therapy, endobronchial stents, and/or brachytherapy.

In other specific embodiments, patients with non-small lung cell cancer are administered a prophylactically or therapeutically effective amount of a TLR3 agonist in combination with the administration of a prophylactically or therapeutically effective amount of one or more other therapies useful for lung cancer treatment or management including but not limited to: palliative radiation therapy, the combination of cisplatin, vinblastine and mitomycin, the combination of cisplatin and vinorelbine, paclitaxel, docetaxel or gemcitabine, the combination of carboplatin and paclitaxel, interstitial radiation therapy for endobronchial lesions or stereotactic radiosurgery

Combination with chemotherapy, preferred examples

An adjunct therapy contemplated in the present invention is chemotherapy. Adjunct chemotherapies may include, for example, cisplatin (CDDP), carboplatin, oxaliplatin, procarbazine, mechlorethamine, cyclophosphamide, camptothecin, ifosfamide, melphalan, chlorambucil, busulfan, nitrosurea, dactinomycin, daunorubicin, doxorubicin, bleomycin, plicomycin, mitomycin, etoposide (VP16), tamoxifen, raloxifene, estrogen receptor binding agents, taxol, gemcitabine, navelbine, farnesyl-protein transferase inhibitors, transplatinum, 5-fluorouracil, vincristin, vinblastin and methotrexate, or any analog or derivative variant of the foregoing.

Taxol/Paclitaxel. Paclitaxel, also known as taxol is a diterpene alkaloid thus it possesses a taxane skeleton in its structure. Paclitaxel is extracted from the bark of the Pacific yew (*Taxus brevifolia*) as a natural compound having anti-cancer activity (Fuchs and Johnson, 1978). Paclitaxel works against cancer by interfering with mitosis. Paclitaxel is a taxoid drug, widely used as an effective treatment of primary and metastatic cancers.

Paclitaxel (Taxol) is widely used in the treatment of breast, ovarian, and other solid tumors. Randomized clinical trials have shown a survival advantage among patients with primary breast cancer who received paclitaxel in addition to anthracycline-containing adjuvant chemotherapy (Eifel et al., 2001). Furthermore, paclitaxel is effective for both metastatic breast cancer (Holmes et al., 1991; Nabholz et al., 1996; Bishop et al., 1999) and advanced ovarian cancer (McGuire et al., 1996; Piccart et al., 2000). The antitumor activity of paclitaxel is unique because it promotes microtubule assembly and stabilizes the microtubules, thus preventing mitosis (Huizing et al.,

1995). Paclitaxel does this by reversibly and specifically binding to the B subunit of tubulin, forming microtubule polymers thereby stabilizing them against depolymerization and thus leading to growth arrest in the G2/M phase of the cell cycle (Gotaskie and Andreassi, 1994). This makes taxol unique in comparison to vincristine and vinblastine which cause microtubule disassembly
5 (Gatzenicier et al., 1995). Additionally, recent evidence indicates that the microtubule system is essential to the release of various cytokines and modulation of cytokine release may play a major role in the drug's antitumor activity (Smith et al., 1995).

However, some patients are resistant to paclitaxel therapy, and the characteristics of patients who will benefit from the drug have not been well defined. Identification of molecular characteristics
10 predictive of paclitaxel sensitivity or resistance could aid in selecting patients to receive this therapy. Thus, in particular embodiments, the present invention relates to paclitaxel sensitivity in a patient having cancer. Previous reports have demonstrated that paclitaxel resistance is due to a variety of mechanisms such as up-regulation of anti-apoptotic Bcl-2 family members, such as Bcl-2 and Bcl-XL (Tang et al., 1994); up-regulation of membrane transporters (e.g., mdr-1), resulting in
15 an increased drug efflux- (Huang et al., 1997); mutations in beta-tubulin resulting in abolishment of paclitaxel binding (Giannakakou et al., 1997); and up-regulation of ErbB2 (HER2) through inhibition of cyclin-dependent kinase-1 (Cdk1), resulting in delayed mitosis (Yu et al., 1998). Due to the antimitotic activity of paclitaxel it is a useful cytotoxic drug in treating several classic refractory tumors. Paclitaxel has primarily been used to treat breast cancer and ovarian cancer. It
20 may also be used in treating head and neck cancer, Kaposi's sarcoma and lung cancer, small cell and non-small cell lung cancer. It may also slow the course of melanoma. Response rates to taxol treatment varies among cancers. Advanced drug refractory ovarian cancer is reported to respond at a 19-36% rate, previously treated metastatic breast cancer at 27-62%, and various lung cancers at 21-37%. Taxol has also been shown to produce complete tumor remission in some cases
25 (Guchelaar et al., 1994).

Paclitaxel is given intravenously since it irritates skin and mucous membranes on contact. It is typically administered intravenously by a 3 to 24 hour infusion three times per week (Guchelaar et al., 1994).

Doxorubicin. Doxorubicin hydrochloride, 5,12-Naphthacenedione, (8s-ds)-10-[(3-amino-
30 2,3,6-trideoxy- α -L-lyxo-hexopyranosyl)oxy]-7,8,9,10-tetrahydro-6,8,11-trihydroxy-g-(hydroxyacetyl)-l-methoxy-hydrochloride (hydroxydaunorubicin hydrochloride, Adriamycin) is used in a wide antineoplastic spectrum. It binds to DNA and inhibits nucleic acid synthesis and mitosis, and promotes chromosomal aberrations.

Administered alone, it is the drug of first choice for the treatment of thyroid adenoma and primary
35 hepatocellular carcinoma. It is a component of first-choice in combination with other agents for the treatment of ovarian tumors, endometrial and breast tumors, bronchogenic oat-cell carcinoma, non-small cell lung carcinoma, gastric adenocarcinoma, retinoblastoma, neuroblastoma, mycosis

fungoides, pancreatic carcinoma, prostatic carcinoma, bladder carcinoma, myelonia, diffuse histiocytic lymphoma, Wilms' tumor, Hodgkin's disease, adrenal tumors, osteogenic sarcoma soft tissue sarcoma, Ewing's sarcoma, rhabdomyosarcoma and acute lymphocytic leukemia. It is an alternative drug for the treatment of islet cell, cervical, testicular and adrenocortical cancers. It is also an immunosuppressant.

Since doxorubicin is poorly absorbed it is administered intravenously. The pharmacokinetics of this chemotherapeutic agent are multicompartamental. Distribution phases have half-lives of 12 minutes and 3.3 hrs. The elimination half-life is about 30 hrs. Forty to 50% is secreted into the bile. Most of the remainder is metabolized in the liver, partly to an active metabolite (doxorubicinol), but a few percent is excreted into the urine. In the presence of liver impairment, the dose should be reduced. Appropriate doses are, for an adult, administered intravenously, are 60 to 75 mg/m² at 2 1-day intervals, or 25 to 30 mg/m² on each of 2 or 3 successive days repeated at 3- or 4-wk intervals, or 20 mg/m² once a week. The lowest dose should be used in elderly patients, when there is prior bone-marrow depression caused by prior chemotherapy or neoplastic marrow invasion, or when the drug is combined with other myelopoietic suppressant drugs. The dose should be reduced by 50% if the serum bilirubin lies between 1.2 and 3 mg/dL and by 75% if above 3 mg/dL. The lifetime total dose should not exceed 550 mg/m² in patients with normal heart function and 400 mg/m² in persons having received mediastinal irradiation. Alternatively, 30 mg/m² on each of 3 consecutive days, repeated every 4 wk may be administered. Exemplary doses may be 10 mg/m², 20 mg/m², 30 mg/m², 50 mg/m², 100 mg/m², 150 mg/m², 175 mg/m², 200 mg/m², 225 mg/m², 250 mg/m², 275 mg/m², 300 mg/m², 350 mg/m², 400 mg/m², 425 mg/m², 450 mg/m², 475 mg/m², 500 mg/m². Of course, all of these dosages are exemplary, and any dosage in-between these points is also expected to be of use in the present invention.

Combination with Radiotherapy

Radiotherapy, also called radiation therapy, involves the use of ionizing radiation to treat cancers and other diseases. Ionizing radiation deposits energy that injures or destroys cells in the area being treated (the "target tissue") by damaging their genetic material, and thereby inhibiting cell proliferation. Ionizing radiation induces the formation of hydroxyl radicals, placing the cells under oxidative stress. These radicals damage DNA, which causes cytotoxicity.

Radiotherapeutic agents that cause DNA damage are well known in the art and have been extensively used. Radiotherapeutic agents, through the production of oxygen-related free radicals and DNA damage, may lead to cell death or apoptosis. These agents may include, but are not limited to, γ -rays, X-rays, and/or the directed delivery of radioisotopes to tumor cells (known as internal radiotherapy). Internal radiotherapy may further include but is not limited to, brachytherapy, interstitial irradiation, and intracavitary irradiation. Other radiotherapeutic agents that are DNA damaging factors include microwaves and UV-irradiation. These factors effect a

broad range of damage on DNA, on the precursors of DNA, on the replication and repair of DNA, and on the assembly and maintenance of chromosomes. Other approaches to radiation therapy are also contemplated in the present invention.

Such techniques may comprise intraoperative irradiation, in which a large dose of external

5 radiation is directed at the tumor and surrounding tissue during surgery; and particle beam radiation therapy which involves the use of fast-moving subatomic particles to treat localized cancers.

Radiotherapy may further involve the use of radiosensitizers and/or radioprotectors to increase the effectiveness of radiation therapy. Radiolabeled antibodies may also be used to deliver doses of radiation directly to the cancer site, this is known as radioimmunotherapy.

10 Dosage ranges for X-rays range from daily doses of 50 to 200 roentgens for prolonged periods of time (3 to 4 wk), to single doses of 2000 to 6000 roentgens. Dosage ranges for radioisotopes vary widely, and depend on the half-life of the isotope, the strength and type of radiation emitted, and the uptake by the neoplastic cells.

Surgery

15 Approximately 60% of persons with cancer will undergo surgery of some type, which includes preventative, diagnostic or staging, curative and palliative surgery. Curative surgery includes resection in which all or part of cancerous tissue is physically removed, excised, and/or destroyed. Tumor resection refers to physical removal of at least part of a tumor. In addition to tumor resection, treatment by surgery includes laser surgery, cryosurgery, electrosurgery, and

20 microscopically controlled surgery. It is further contemplated that the present invention may be used in conjunction with removal of superficial cancers, precancers, or incidental amounts of normal tissue.

Upon excision of part of all of cancerous cells, tissue, or tumor, a cavity may be formed in the body. Treatment may be accomplished by administration of an additional anti-cancer therapy, more

25 particularly a TLR3 agonist. Such treatment may be repeated, for example, every 1, 2, 3, 4, 5, 6, or 7 days, or every 1, 2, 3, 4, and 5 weeks or every 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12 months. These treatments may be of varying dosages as well.

Hormonal Therapy

Hormonal therapy may also be used in conjunction with the present invention or in combination

30 with any other cancer therapy previously described. The use of hormones may be employed in the treatment of certain cancers such as breast, prostate, ovarian, or cervical cancer to lower the level or block the effects of certain hormones such as testosterone or estrogen. This treatment is often used in combination with at least one other cancer therapy as a treatment option or to reduce the risk of metastases.

35 *Other agents*

It is contemplated that other agents may be used in combination with the present invention to improve the therapeutic efficacy of treatment. These additional agents include immunomodulatory

agents, agents that affect the upregulation of cell surface receptors and GAP junctions, cytostatic and differentiation agents, or inhibitors of cell adhesion. Immunomodulatory agents include tumor necrosis factor; interferon alpha, beta, and gamma; *JL-2*, IL-12, IL-15, IL-21 and other cytokines ; F42K and other cytokine analogs; or MIP- 1, MIP- 1beta, MCP- 1, RANTES, and other

5 chemokines.

The treatment with a TLR3 agonist, and more particularly dsRNA molecule, may be accomplished as disclosed in the literature cited above. Furthermore, the treatment may be performed either alone or in combination with other drugs or treatments. The treatment may include a reduction in tumor size, a reduction or delay in tumor growth, development or metastasis, or a regression of cancer.

10 Further aspects and advantages of this invention will be disclosed in the following examples, which should be regarded as illustrative and not limiting the scope of this invention.

EXAMPLES

Toll like receptor 3 (TLR3) is known to be expressed by myeloid dendritic cells (DC) and to induce

15 their maturation following binding with double stranded RNA (dsRNA) or its synthetic homologues polyAU and polyLC. Several clinical trials have reported that injection of dsRNA is associated with survival benefit in cancer patients. In the present study, the inventors have asked whether dsRNA could act directly on tumor cells through TLR3. Patients and methods : 300

patients with early breast cancer have been included from 1972 to 1979 in a randomized trial

20 comparing post-operative administration of polyAU with no treatment. Results have been reported that showed a trend for a survival benefit in patients with involved axillary lymph nodes (n=200). , Tumor biopsies from these patients were stained with TLR3-specific mAb and correlation between TLR3 expression and polyAU efficacy was determined.

To investigate directly the effects of dsRNA, both freshly isolated breast tumor cells and cancer

25 cell lines were cultured with polyI:C, and apoptosis was measured. The involvement of TLR3 in cell response was established by TLR3 RNA interference.

Results : 182 tumor samples (91%) were available from the 200 pTxN+MO patients included in this randomized trial. TLR3 was strongly expressed by tumor cells in 18 patients (10%). Table 1 reports the 20-year survival rates according to treatment and TLR3 expression.

30

Targeting Toll like receptor 3 in breast cancer : results of randomized trial and in vitro studies

Material and methods:

Patients:

35 200 patients were included in the present study. All patients had been previously included in a prospective randomized trial that compared double stranded RNA (polyAU) to placebo. This trial have already been reported elsewhere. Briefly, this randomized trial included patients with Tl-

3N0-3M0 breast cancer treated with surgery. Treatment consisted in weekly iv injection of polyAU (Beaufour Ipsen). A total of 6 injections were performed. PolyAU was administered at a fixed dose of 60 mg/injection. This trial initially included 300 patients. Since initial results of the trial reported a trend for benefit only in patients with axillary lymph node involvement, only the 200 patients with axillary node involvement were included in the present study.

Immunostainings:

Tumor blocks were available in 182 out of 200 patients included in the present study. Paraffin-embedded, 5 um-thick tissue section from all 182 tumors were stained with either polyclonal antiTLR3 (gift from Dr Pobolsky, Massachusetts General Hospital, Boston) or rabbit preimtnun serum. A mouse monoclonal anti-rabbit IgG was used as secondary antibody. Immunostainings were assessed by 2 pathologists who were blinded for clinical files. The TLR3 expression was classified according to the percentage of tumor cells stained and the intensity of staining. A tumor was classified as positive when more than 10% of tumor cells were strongly stained with the anti-TLR3 antibody.

Statistics:

Survival curves were determined according to Kaplan-Meier method. Survival curves were compared using Khi2 test.

Results:

Patients characteristics

One hundred eighty two tumors were processed. The immunostaining could not be interpreted in 7 patients (absence of tumor cells in 4 patients, artefact in 3 patients). The analysis was therefore performed on 175 patients. This represents 87% of the patients included in the randomized trial. The median follow-up of living patients was 23 years (12 to 26 years). The patients characteristics are reported in Table 1. Briefly, the median age is 50, the median number number of lymph node involved was 4 (1-31), 26% of tumor were staged pT3 and 35% were classified as grade IH according to Scarf and Bloom Richardson.

Table 1 : Patients characteristics

Characteristics	TLR3- tumors (n=157)		TLR3+ tumors (n=18)		Total (n=175)
	Observation (n=77)	Poly AU (n=80)	Observation (n=10)	Poly AU (n=8)	
Age (median)	50	50	52	49	50
Nb lymph node involved (median)	5 (1-31)	4 (1-27)	2 (1-8)	4 (1-9)	4 (1-31)
pT					
pT1	8	1	0	0	9
pT2	56	54	6	6	122
pT3	13	27	4	2	46

Tumor grade					
I	11	9	1	2	23
II	34	49	6	3	92
III	32	24	3	3	62
Post-operative radiotherapy?					
Yes	74	77	9	8	168
No	3	3	1	0	7

Immunostainings

TLR3 was strongly expressed by tumor cells in 18 samples (10.4% of assessable tumors).

Immunostainings are shown in Figure 1. TLR3 was mainly expressed on the cell surface and

5 cytoplasm of tumor cells. In situ carcinoma and normal breast tissues were stained by anti-TLR3 in most cases. The patients characteristics of the TLR3+ tumors did not differ to that of TLR3- tumors (Table 1).

Correlation between TLR3 expression and survival after treatment with polyAU

10 The 20 year OS of patients treated or not with polyAU were 42% and 35% respectively (p=0.09). When only patients with TLR3- tumors were considered, the 20 year OS were 41% for patients treated with polyAU, and 37% for those assigned to observation arm (p=0.52) (Figure 2a). When only patients presenting TLR3+ tumors were considered, the 20 year OS were 88% for patients treated with polyAU, and 22% for patients assigned to the observation arm (p=0.01) (Figure 2b).

15

Conclusion:

TLR3 is overexpressed by tumor cells in around 10% of cancer cases.

TLR3 expression correlates with the benefit of adjuvant therapy with polyAU in patients with lymph node positive breast cancer.

20

CLAIMS

What is claimed is:

- 5 L A method for treating a subject having a cancer, the method comprising the steps of:
- a. determining whether cells of said cancer in said subject express a TLR3 receptor, the expression of a TLR3 receptor being indicative of a subject responding to a TLR3 agonist, and
 - b. administering to said subject whose cancer cells are determined to express a
- 10 TLR3 receptor a pharmaceutical composition comprising an effective amount of a TLR3 agonist and a pharmaceutically acceptable carrier.
2. The method according to claim 1, wherein said TLR3 agonist is a dsRNA TLR3 agonist.
- 15 3. The method according to claim 2, wherein said dsRNA TLR3 agonist is at least 90% double-stranded.
4. The method according to claim 2 or 3, wherein said dsRNA TLR3 agonist is selected
- 20 from polyA:polyU, polyLpolyC, or polyI:poly(Ci₂U).
5. The method according to claim 4, wherein said dsRNA is a fully double stranded polyA:polyU molecule consisting of between 19 and 30 base pairs and comprising between 1 and 30 stabilizing modifications, or a 3' cap, or a 5' cap.
- 25 6. The method according to any one of claims 1-5, wherein said cancer is selected from a solid tumor or a carcinoma.
7. The method of claim 6, wherein the solid tumor is selected from breast cancer, colon
- 30 cancer, lung cancer, renal cancer, metastatic or invasive malignant melanoma, prostate cancer, brain tumor, bladder cancer and liver cancer.
8. The method according to claim 7, wherein said solid tumor is a breast cancer.
- 35 9. The method according to claim 8, wherein said breast cancer is a lymph node-positive breast cancer involving 1 to 3 lymph nodes.

10. The method according to anyone of claims 1-9, wherein said determining whether cells of said cancer in said subject express a TLR3 receptor is performed ex vivo on a biopsied sample of said cancer using a TLR3-specific ligand.

5 11. The method according to claim 10, wherein said TLR3-specific ligand is an antibody or a fragment or a derivative thereof.

12. The method according to claim 10, wherein said TLR3-specific ligand is a TLR3-specific nucleic acid primer or probe.

10

13. The method according to claim 6 comprising the additional step of surgically removing a portion of said solid tumor prior to administering said TLR3 agonist.

14. The method according to anyone of claims 1-13, wherein said subject is administered
15 said TLR3 agonist about once every week for at least about six weeks.

15. The method according to any one of claims 1-14, comprising the additional step of administering to said subject a second therapeutic agent selected from a TLR3 agonist of different molecular composition than the TLR3 agonist administered in step b; a cytotoxin, a cytotoxin-
20 TLR3 ligand complex, an agent that inhibits expression or an activity of a tumor antigen, an agent that inhibits expression or an activity of a tumor proliferative protein, a chemotherapy agent, an agent or combination of agents typically used for the treatment of the specific cancer to be treated, an immunostimulatory agent, an immunosuppression agent, a cytokine or cytokine analog, a chemokines, an agent that affects upregulation of a cell surface receptor, an agent that affects GAP
25 junctions, a cytostatic agent, an agent that inhibits differentiation, or an agent that inhibits cell adhesion, wherein said second therapeutic agent is administered separately as part of a multiple dosage regimen, or as part of said pharmaceutical composition.

16. The method according to anyone of claims 1-16, additionally comprising the step of
30 treating said subject with radiotherapy.

17. The method according to claim 13, wherein said TLR3 agonist composition is formulated into an implantable drug release device that is implanted near the site of said solid tumor during said surgical procedure.

35

18. The method according to claim 3, wherein said dsTLR3 agonist is an siRNA or an shRNA that specifically binds to mRNA encoding a tumor antigen or another protein involved in

tumor proliferation.

19. A method for treating a subject having a cancer, the method comprising the steps of:
- a. administering to said subject an effective amount of an agent that causes
 - 5 upregulation of TLR3 expression, and
 - b. administering to said subject an effective amount of a TLR3 agonist.

20. The method according to claim 19, comprising the additional step of determining whether cells of said cancer in said subject express a TLR3 receptor, said additional step being
- 10 performed prior to step b.

21. A composition of matter for treating a subject having a cancer comprising, in separate dosage forms:
- a. an agent that causes upregulation of TLR3 expression, and
 - 15 b. a TLR3 agonist,
- wherein said agent and said agonist are associated with one another in said composition of matter.

22. A kit for determining if an agent is capable of increasing the susceptibility of a cancer to treatment with a TLR3 agonist, wherein the kit comprises:
- 20 a. a cancer cell or cancer cell line which can express TLR3;
 - b. a TLR3-specific ligand; and
 - c. a reagent for detecting the binding of said TLR3-specific ligand to a TLR3 expressed by said cancer cell.

23. A pharmaceutical composition comprising:
- a. a TLR3 agonist;
 - b. a second therapeutic agent selected from a TLR3 agonist of different molecular composition the TLR3 agonist of part a; a cytotoxin, a cytotoxin-TLR3 ligand complex, an agent that inhibits expression or an activity of a tumor antigen, an agent that inhibits
 - 30 expression or an activity of a tumor proliferative protein, a chemotherapy agent, an agent or combination of agents typically used for the treatment of the specific cancer to be treated, an immunostimulatory agent, an immunosuppression agent, a cytokine or cytokine analog, a chemokines, an agent that affects upregulation of a cell surface receptor, an agent that affects GAP junctions, a cytostatic agent, an agent that inhibits differentiation, an
 - 35 agent that inhibits cell adhesion, or an endocytosis inhibitor; and
 - c. a pharmaceutically acceptable carrier.

24. A composition of matter for treating a subject having a cancer comprising, in separate dosage forms:

- a. a TLR3 agonist; and
- b. a second therapeutic agent selected from a TLR3 agonist of different molecular composition the TLR3 agonist of part a; a cytotoxin, a cytotoxin-TLR3 ligand complex, an agent that inhibits expression or an activity of a tumor antigen, an agent that inhibits expression or an activity of a tumor proliferative protein, a chemotherapy agent, an agent or combination of agents typically used for the treatment of the specific cancer to be treated, an immunostimulatory agent, a TLR2, TLR4, TLR7, TLR8 or TLR9 agonist, an immunosuppression agent, a cytokine or cytokine analog, a chemokines, an agent that affects upregulation of a cell surface receptor, an agent that affects GAP junctions, a cytostatic agent, a pro-apoptotic agent, an agent that inhibits differentiation, an agent that inhibits cell adhesion, or an endocytosis inhibitor, wherein said agent and said agonist are associated with one another in said composition of matter.

25. A method of impregnating or filling an implantable drug release device comprising the step of contacting said drug release device with a TLR3 agonist or a composition comprising a TLR3 agonist.

26. An implantable drug release device impregnated with or containing a TLR3 agonist or a composition comprising a TLR3 agonist, such that said TLR3 agonist is released from said device and is therapeutically active.

27. A method for assessing the response of a subject having cancer to a treatment using a TLR3 agonist or selecting a subject having a cancer that responds to a treatment using a TLR3 agonist, the method comprising the step of determining whether cancer cells in said subject express a TLR3 receptor, the expression of a TLR3 receptor being indicative of a responder subject.

28. A kit for assessing the response of a subject having cancer to a treatment using a TLR3 agonist or for selecting subjects having a cancer that responds to a treatment using a TLR3 agonist, the kit comprising:

- a. a TLR3-specific ligand; and
- b. additional reagents for detecting the expression of a TLR3 receptor in a cancer cell in a sample.

29. The kit of claim 28, wherein the TLR3-specific ligand selected from an antibody or a fragment or a derivative thereof; a TLR3-specific nucleic acid primer or probe, or a double-stranded RNA molecule.

5 30. A complex comprising:

- a. a TLR3 ligand; and
- b. a cytotoxic agent,

wherein said agonist and said cytotoxic agent are bound to one another directly through a covalent bond, bond to one another through a non-covalent bond, conjugated to one another directly or
10 through a linking moiety or are present in a single pharmaceutical carrier or vehicle.

31. The complex according to claim 30, wherein said TLR3 ligand is selected from an antibody, or a fragment or derivative thereof, that specifically binds to TLR3; or a double-stranded RNA molecule.

15

32. A method of determining if a test compound is useful for the treatment of cancer, said method comprising the step of determining if said test compound is a TLR3 agonist.

33. The method according to claim 32, wherein said determining if a test compound is
20 useful for the treatment of cancer comprises the steps of:

- a. contacting said test compound with a cancer cell characterized by the presence of TLR3 on its cell surface; and
- b. determining if said test compound induces a TLR3-mediated biological effect.

25 34. The method according to claim 33, wherein said TLR3-mediated biological effect is apoptosis or cell death of said cancer cell.

35. A kit for determining if a test compound is useful for the treatment of cancer, said kit comprising:

- 30 a. a cancer cell characterized by the presence of TLR3 on its cell surface; and
- b. a reagent for detecting if said test compound causes TLR3-mediated apoptosis or cell death of said cancer cell; and
- optionally comprising a reagent that detects agonism of TLR3 by said test compound.

35 36. A use of a TLR3 agonist for the manufacture of a medicament for treating cancer in a subject, wherein said cancer in said subject comprises cancer cells expressing a TLR3 receptor.

37. The use according to claim 36, wherein said TLR3 agonist is a dsRNA TLR3 agonist.

38. The use according to claim 37, wherein said dsRNA TLR3 agonist is at least 90% double-stranded.

5

39. The use according to claim 38, wherein said dsRNA TLR3 agonist is selected from polyA:polyU, polyI:polyC, or polyI:poly(C12U).

10 40. The use according to claim 39, wherein said dsRNA is a fully double stranded polyA:polyU molecule consisting of between 19 and 30 base pairs and comprising between 1 and 30 stabilizing modifications, or a 3' cap, or a 5' cap.

41. The use according to anyone of claims 36-40, wherein said cancer is selected from a solid tumor or a carcinoma.

15

42. The use of claim 41, wherein the solid tumor is selected from breast cancer, colon cancer, lung cancer, renal cancer, metastatic or invasive malignant melanoma, prostate cancer, brain tumor, ladder cancer and liver cancer.

20

43. The use according to claim 42, wherein said solid tumor is a breast cancer.

44. The use according to claim 43, wherein said breast cancer is a lymph node-positive breast cancer involving 1 to 3 lymph nodes.

25

45. The use according to anyone of claims 36-44, wherein a determination that said cancer expresses a TLR3 receptor is achieved by using contacting a TLR3-specific ligand ex vivo with a biopsied sample of said cancer.

30

46. The use according to claim 45, wherein said TLR3-specific ligand is an antibody or a fragment or a derivative thereof.

47. The use according to claim 46, wherein said TLR3-specific ligand is a TLR3-specific nucleic acid primer or probe.

35

48. The use according to claim 41, wherein said medicament is indicated for use in a subject following surgical removal of a portion of said solid tumor.

49. The use according to anyone of claims 3648, wherein said medicament is indicated for administration to said patient about once every week for at least about six weeks.


1/3

Figure 1




2/3

Figure 2a: Survival of patients with TLR3- tumors according to treatment



3/3

Figure 2b: Survival of patients with TLR3+ tumors according to treatment



1
SEQUENCE LISTING

<110> Institut Gustave Roussy

<120> IMPROVED TREATMENT OF CANCER USING TLR3 AGONISTS

<130> B0331WO2

<150> PCT/IB2004/004093

<151> 2004-11-19

<150> US 11/144,322

<151> 2005-06-03

<160> 4

<170> Patentln version 3.1

<210> 1

<211> 3057

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (102) ..(2816)

<223>

<400> 1

cacttttcgag agtgccgtct atttgccaca cacttccctg atgaaatgtc tggatttgga 60

ctaaagaaaa aaggaaaggc tagcagtcac ccaacagaat c atg aga cag act ttg 116
Met Arg Gin Thr Leu
1 5

cct tgt ate tac ttt tgg ggg ggc ctt ttg ccc ttt ggg atg ctg tgt 164
Pro Cys lie Tyr Phe Trp Gly Gly Leu Leu Pro Phe Gly Met Leu Cys
10 15 20

gca tec tec ace ace aag tgc act gtt age cat gaa gtt get gac tgc 212
Ala Ser Ser Thr Thr Lys Cys Thr Val Ser His Glu Val Ala Asp Cys
25 30 35

age cac ctg aag ttg act cag gta ccc gat gat eta ccc aca aac ata 260
ser His Leu Lys Leu Thr Gin Val Pro Asp Asp Leu Pro Thr Asn lie
40 45 50

aca gtg ttg aac ctt ace cat aat caa etc aga aga tta cca gee gee 308
Thr Val Leu Asn Leu Thr His Asn Gin Leu Arg Arg Leu Pro Ala Ala
55 60 65

aac ttc aca agg tat age cag eta act age ttg gat gta gga ttt aac 356
Asn Phe Thr Arg Tyr Ser Gin Leu Thr Ser Leu Asp Val Gly Phe Asn
70 75 80 85

ace ate tea aaa ctg gag cca gaa ttg tgc cag aaa ctt ccc atg tta 404
Thr lie Ser Lys Leu Glu Pro Glu Leu Cys Gin Lys Leu Pro Met Leu
90 95 100

aaa gtt ttg aac etc cag cac aat gag eta tct caa ctt tct gat aaa 452
Lys Val Leu Asn Leu Gin His Asn Glu Leu Ser Gin Leu Ser Asp Lys
105 110 115

ace ttt gee ttc tgc acg aat ttg act gaa etc cat etc atg tec aac 500
Thr Phe Ala Phe cys Thr Asn Leu Thr Glu Leu His Leu Met Ser Asn
120 125 130

tea ate cag aaa att aaa aat aat ccc ttt gtc aag cag aag aat tta 548

2

Ser	lie	Gin	Lys	lie	Lys	Asn	Asn	pro	Phe	VaI	Lys	Gin	Lys	Asn	Leu	
	135					140					145					
ate	aca	tta	gat	ctg	tct	cat	aat	ggc	ttg	tea	tct	aca	aaa	tta	gga	596
lie	Thr	Leu	Asp	Leu	ser	His	Asn	Gly	Leu	Ser	ser	Thr	Lys	Leu	GIy	
	150				155					160					165	
act	cag	gtt	cag	ctg	gaa	aat	etc	caa	gag	ctt	eta	tta	tea	aac	aat	644
Thr	Gin	VaI	Gin	Leu	Glu	Asn	Leu	Gin	Glu	Leu	Leu	Leu	Ser	Asn	Asn	
				170					175					180		
aaa	att	caa	gcg	eta	aaa	agt	gaa	gaa	ctg	gat	ate	ttt	gcc	aat	tea	692
Lys	lie	Gin	Ala	Leu	Lys	ser	Glu	Glu	Leu	Asp	lie	Phe	Ala	Asn	Ser	
			185					190					195			
tct	tta	aaa	aaa	tta	gag	ttg	tea	teg	aat	caa	att	aaa	gag	ttt	tct	740
Ser	Leu	Lys	Lys	Leu	Glu	Leu	Ser	Ser	Asn	Gin	lie	Lys	Glu	Phe	ser	
		200					205					210				
cca	ggg	tgt	ttt	cac	gca	att	gga	aga	tta	ttt	ggc	etc	ttt	ctg	aac	788
Pro	Gly	Cys	Phe	His	Ala	lie	GIy	Arg	Leu	Phe	Gly	Leu	Phe	Leu	Asn	
	215					220					225					
aat	gtc	cag	ctg	ggt	ccc	age	ctt	aca	gag	aag	eta	tgt	ttg	gaa	tta	836
Asn	VaI	Gin	Leu	Gly	Pro	ser	Leu	Thr	Glu	Lys	Leu	Cys	Leu	Glu	Leu	
	230				235					240				245		
gca	aac	aca	age	att	egg	aat	ctg	tct	ctg	agt	aac	age	cag	ctg	tec	884
Ala	Asn	Thr	ser	lie	Arg	Asn	Leu	Ser	Leu	Ser	Asn	Ser	Gin	Leu	Ser	
				250					255					260		
ace	ace	age	aat	aca	act	ttc	ttg	gga	eta	aag	tgg	aca	aat	etc	act	932
Thr	Thr	ser	Asn	Thr	Thr	Phe	Leu	Gly	Leu	Lys	Trp	Thr	Asn	Leu	Thr	
			265					270					275			
atg	etc	gat	ctt	tec	tac	aac	aac	tta	aat	gtg	gtt	ggt	aac	gat	tec	980
Met	Leu	Asp	Leu	Ser	Tyr	Asn	Asn	Leu	Asn	VaI	val	Gly	Asn	Asp	Ser	
		280					285					290				
ttt	get	tgg	ctt	cca	caa	eta	gaa	tat	ttc	ttc	eta	gag	tat	aat	aat	1028
Phe	Ala	Trp	Leu	Pro	Gin	Leu	Glu	Tyr	Phe	Phe	Leu	Glu	Tyr	Asn	Asn	
	295					300					305					
ata	cag	cat	ttg	ttt	tct	cac	tct	ttg	cac	ggg	ctt	ttc	aat	gtg	agg	1076
lie	Gin	His	Leu	Phe	Ser	His	Ser	Leu	His	Gly	Leu	Phe	Asn	VaI	Arg	
	310				315					320					325	
tac	ctg	aat	ttg	aaa	egg	tct	ttt	act	aaa	caa	agt	att	tec	ctt	gcc	1124
Tyr	Leu	Asn	Leu	Lys	Arg	ser	Phe	Thr	Lys	Gin	Ser	lie	ser	Leu	Ala	
				330					335					340		
tea	etc	ccc	aag	att	gat	gat	ttt	tct	ttt	cag	tgg	eta	aaa	tgt	ttg	1172
Ser	Leu	Pro	Lys	lie	Asp	Asp	Phe	Ser	Phe	Gin	Trp	Leu	Lys	Cys	Leu	
			345				350						355			
gag	cac	ctt	aac	atg	gaa	gat	aat	gat	att	cca	ggc	ata	aaa	age	aat	1220
Glu	His	Leu	Asn	Met	Glu	Asp	Asn	Asp	lie	Pro	Gly	lie	Lys	Ser	Asn	
			360				365					370				
atg	ttc	aca	gga	ttg	ata	aac	ctg	aaa	tac	tta	agt	eta	tec	aac	tec	1268
Met	Phe	Thr	Gly	Leu	lie	Asn	Leu	Lys	Tyr	Leu	Ser	Leu	ser	Asn	Ser	
						380					385					
ttt	aca	agt	ttg	cga	act	ttg	aca	aat	gaa	aca	ttt	gta	tea	ctt	get	1316
Phe	Thr	ser	Leu	Arg	Thr	Leu	Thr	Asn	Glu	Thr	Phe	VaI	ser	Leu	Ala	
					395				400						405	
cat	tct	ccc	tta	cac	ata	etc	aac	eta	ace	aag	aat	aaa	ate	tea	aaa	1364
His	Ser	Pro	Leu	His	lie	Leu	Asn	Leu	Thr	Lys	Asn	Lys	H e	Ser	Lys	

3																
410						415						420				
ata lie	gag Glu	agt Ser	gat Asp 425	get Ala	ttc Phe	tct Ser	tgg Trp	ttg Leu 430	ggc Gly	cac His	eta Leu	gaa Glu 435	gta Val	ctt Leu	gac Asp	1412
ctg Leu	ggc Gly	ctt Leu 440	aat Asn	gaa Glu	att lie	ggg Gly	caa Gin 445	gaa Glu	etc Leu	aca Thr	ggc Gly	cag Gin 450	gaa Glu	tgg Trp	aga Arg	1460
ggc Gly	eta Leu 455	gaa Glu	aat Asn	att lie	ttc Phe	gaa Glu 460	ate lie	tat Tyr	ctt Leu	tec Ser	tac Tyr 465	aac Asn	aag Lys	tac Tyr	ctg Leu	1508
cag Gin 470	ctg Leu	act Thr	agg Arg	aac Asn	tec Ser 475	ttt Phe	gee Ala	ttg Leu	gtc Val	cca Pro 480	age Ser	ctt Leu	caa Gin	cga Arg	ctg Leu 485	1556
atg Met	etc Leu	cga Arg	agg Arg	gtg Val 490	gee Ala	ctt Leu	aaa Lys	aat Asn	gtg Val 495	gat Asp	age Ser	tct ser	cct Pro	tea Ser 500	cca Pro	1604
ttc Phe	cag Gin	cct Pro	ctt Leu 505	cgt Arg	aac Asn	ttg Leu	ace Thr	att lie 510	ctg Leu	gat Asp	eta Leu	age Ser	aac Asn 515	aac Asn	aac Asn	1652
ata lie	gcc Ala	aac Asn 520	ata lie	aat Asn	gat Asp	gac Asp	atg Met 525	ttg Leu	gag Glu	ggc Gly	ctt Leu	gag Glu 530	aaa Lys	eta Leu	gaa Glu	1700
att lie	etc Leu 535	gat Asp	ttg Leu	cag Gin	cat His	aac Asn 540	aac Asn	tta Leu	gca Ala	egg Arg	etc Leu 545	tgg Trp	aaa Lys	cac His	gca Ala	1748
aac Asn 550	cct Pro	ggc Gly	ggc Gly	ccc Pro	att lie 555	tat Tyr	ttc Phe	eta Leu	aag Lys	ggc Gly 560	ctg Leu	tct Ser	cac His	etc Leu	cac His 565	1796
ate lie	ctt Leu	aac Asn	ttg Leu	gag Glu 570	tec Ser	aac Asn	ggc Gly	ttt Phe	gac Asp 575	gag Glu	ate lie	cca Pro	ggt Val	gag Glu 580	gtc Val	1844
ttc Phe	aag Lys	gat Asp	tta Leu 585	ttt Phe	gaa Glu	eta Leu	aag Lys	ate lie 590	ate lie	gat Asp	tta Leu	gga Gly	ttg Leu 595	aat Asn	aat Asn	1892
tta Leu	aac Asn	aca Thr 600	ctt Leu	cca Pro	gca Ala	tct ser	gtc Val 605	ttt Phe	aat Asn	aat Asn	cag Gin	gtg Val 610	tct Ser	eta Leu	aag Lys	1940
tea Ser	ttg Leu 615	aac Asn	ctt Leu	cag Gin	aag Lys	aat Asn 620	etc Leu	ata lie	aca Thr	tec Ser	ggt Val 625	gag Glu	aag Lys	aag Lys	ggt Val	1988
ttc Phe 630	ggc Gly	cca Pro	get Ala	ttc Phe	agg Arg 635	aac Asn	ctg Leu	act Thr	gag Glu	tta Leu 640	gat Asp	atg Met	cgc Arg	ttt Phe	aat Asn 645	2036
ccc Pro	ttt Phe	gat Asp	tgc cys	acg Thr 650	tgt Cys	gaa Glu	agt Ser	att lie	gcc Ala 655	tgg Trp	ttt Phe	ggt Val	aat Asn	tgg Trp 660	att lie	2084
aac Asn	gag Glu	ace Thr	cat His 665	ace Thr	aac Asn	ate lie	cct Pro	gag Glu 670	ctg Leu	tea ser	age Ser	cac His	tac Tyr 675	ctt Leu	tgc cys	2132
aac Asn	act Thr	cca Pro 680	cct Pro	cac His	tat Tyr	cat His	ggg Gly 685	ttc Phe	cca pro	gtg Val	aga Arg	ctt Leu 690	ttt Phe	gat Asp	aca Thr	2180

4

tea	tct	tgc	aaa	gac	agt	gcc	ccc	ttt	gaa	etc	ttt	ttc	atg	ate	aat	2228
ser	ser	Cys	Lys	Asp	Ser	Ala	Pro	Phe	Glu	Leu	phe	Phe	Met	lie	Asn	
	695					700					705					
ace	agt	ate	ctg	ttg	att	ttt	ate	ttt	att	gta	ctt	etc	ate	cac	ttt	2276
Thr	ser	lie	Leu	Leu	lie	Phe	lie	Phe	lie	VaI	Leu	Leu	lie	His	Phe	
	710				715					720					725	
gag	ggc	tgg	agg	ata	tct	ttt	tat	tgg	aat	gtt	tea	gta	cat	cga	gtt	2324
Glu	GIy	Trp	Arg	lie	Ser	Phe	Tyr	Trp	Asn	VaI	ser	val	His	Arg	VaI	
				730					735					740		
ctt	ggc	ttc	aaa	gaa	ata	gac	aga	cag	aca	gaa	cag	ttt	gaa	tat	gca	2372
Leu	Gly	Phe	Lys	Glu	lie	Asp	Arg	Gin	Thr	Glu	Gin	Phe	Glu	Tyr	Ala	
			745					750					755			
gca	tat	ata	att	cat	gcc	tat	aaa	gat	aag	gat	tgg	gtc	tgg	gaa	cat	2420
Ala	Tyr	lie	lie	His	Ala	Tyr	Lys	Asp	Lys	Asp	Trp	VaI	Trp	Glu	His	
		760					765					770				
ttc	tct	tea	atg	gaa	aag	gaa	gac	caa	tct	etc	aaa	ttt	tgt	ctg	gaa	2468
Phe	ser	Ser	Met	Glu	Lys	Glu	Asp	Gin	Ser	Leu	Lys	Phe	Cys	Leu	Glu	
	775					780					785					
gaa	agg	gac	ttt	gag	gcg	ggc	ggt	ttt	gaa	eta	gaa	gca	att	gtt	aac	2516
Glu	Arg	Asp	Phe	Glu	Ala	Gly	VaI	Phe	Glu	Leu	Glu	Ala	lie	VaI	Asn	
	790				795					800					805	
age	ate	aaa	aga	age	aga	aaa	att	att	ttt	ggt	ata	aca	cac	cat	eta	2564
ser	lie	Lys	Arg	Ser	Arg	Lys	lie	lie	Phe	VaI	lie	Thr	His	His	Leu	
				810					815					820		
tta	aaa	gac	cca	tta	tgc	aaa	aga	ttc	aag	gta	cat	cat	gca	ggt	caa	2612
Leu	Lys	Asp	Pro	Leu	Cys	Lys	Arg	Phe	Lys	VaI	His	His	Ala	VaI	Gin	
			825					830					835			
caa	get	att	gaa	caa	aat	ctg	gat	tec	att	ata	ttg	ggt	ttc	ctt	gag	2660
Gin	Ala	lie	Glu	Gin	Asn	Leu	Asp	Ser	lie	lie	Leu	VaI	Phe	Leu	Glu	
		840					845					850				
gag	att	cca	gat	tat	aaa	ctg	aac	cat	gca	etc	tgt	ttg	cga	aga	gga	2708
Glu	lie	Pro	Asp	Tyr	Lys	Leu	Asn	His	Ala	Leu	Cys	Leu	Arg	Arg	Gly	
	855					860					865					
atg	ttt	aaa	tct	cac	tgc	ate	ttg	aac	tgg	cca	ggt	cag	aaa	gaa	egg	2756
Met	Phe	Lys	Ser	His	Cys	lie	Leu	Asn	Trp	Pro	VaI	Gin	Lys	Glu	Arg	
	870				875				880						885	
ata	ggc	gcc	ttt	cgt	cat	aaa	ttg	caa	gta	gca	ctt	gga	tec	aaa	aac	2804
lie	Gly	Ala	Phe	Arg	His	Lys	Leu	Gin	VaI	Ala	Leu	GIy	ser	Lys	Asn	
				890					895					900		
tct	gta	cat	taa	atttatttaa	atattcaatt	agcaaaggag	aaactttctc									2856
Ser	VaI	His														
aatttaaaaa	ggtctatggc	aaattttaagt	tttccataaa	ggtggtataa	tttggtttatt											2916
catatttgta	aatgattata	ttctatcaca	attacatctc	ttctaggaaa	atgtgtctcc											2976
ttatttcagg	cctatttttg	acaattgact	taattttacc	caaaataaaa	catataagca											3036
cgtaaaaaaa	aaaaaaaaaa	a														3057

<210> 2
 <211> 904
 <212> PRT

5

<213> Homo sapiens

<400> 2

Met 1	Arg	Gln	Thr	Leu 5	pro	Cys	lie	Tyr	Phe 10	Trp	Gly	Gly	Leu	Leu 15	Pro
Phe	Gly	Met	Leu 20	Cys	Ala	Ser	Ser	Thr 25	Thr	Lys	Cys	Thr	Val 30	ser	His
Gl u	Val	Ala 35	Asp	Cys	ser	Hi s	Leu 40	Lys	Leu	Thr	Gln	Val 45	Pro	Asp	Asp
Leu	Pro 50	Thr	Asn	lie	Thr	Val 55	Leu	Asn	Leu	Thr	His 60	Asn	Gln	Leu	Arg
Arg 65	Leu	Pro	Ala	Ala	Asn 70	Phe	Thr	Arg	Tyr	ser 75	Gln	Leu	Thr	Ser	Leu 80
Asp	Val	Gly	Phe	Asn 85	Thr	lie	Ser	Lys	Leu 90	Glu	Pro	Glu	Leu	Cys 95	Gln
Lys	Leu	Pro	Met 100	Leu	Lys	Val	Leu	Asn 105	Leu	Gln	His	Asn	Glu 110	Leu	ser
Gln	Leu	Ser 115	Asp	Lys	Thr	Phe	Ala 120	Phe	Cys	Thr	Asn	Leu 125	Thr	Glu	Leu
His	Leu 130	Met	Ser	Asn	ser	lie 135	Gln	Lys	lie	Lys	Asn 140	Asn	Pro	Phe	Val
Lys 145	Gln	Lys	Asn	Leu	lie 150	Thr	Leu	Asp	Leu	Ser 155	His	Asn	Gly	Leu	ser 160
Ser	Thr	Lys	Leu	Gly 165	Thr	Gln	Val	Gln	Leu 170	Glu	Asn	Leu	Gln	Glu 175	Leu
Leu	Leu	Ser	Asn 180	Asn	Lys	lie	Gln	Ala 185	Leu	Lys	ser	Glu	Glu 190	Leu	Asp
lie	Phe	Ala 195	Asn	ser	ser	Leu	Lys 200	Lys	Leu	Glu	Leu	Ser 205	Ser	Asn	Gln
lie	Lys 210	Glu	Phe	Ser	Pro	Gly 215	Cys	Phe	Hi s	Ala	lie 220	Gly	Arg	Leu	Phe
Gly 225	Leu	Phe	Leu	Asn	Asn 230	Val	Gln	Leu	Gly	Pro 235	ser	Leu	Thr	Glu	Lys 240
Leu	Cys	Leu	Glu	Leu 245	Ala	Asn	Thr	Ser	lie 250	Arg	Asn	Leu	Ser	Leu 255	ser
Asn	Ser	Gln	Leu	Ser	Thr	Thr	Ser	Asn	Thr	Thr	Phe	Leu	Gly	Leu	Lys

6

260						265						270					
Trp	Thr	Asn 275	Leu	Thr	Met	Leu	Asp 280	Leu	Ser	Tyr	Asn	Asn 285	Leu	Asn	val		
val	Gly 290	Asn	Asp	Ser	Phe	Ala 295	Trp	Leu	Pro	Gln	Leu 300	Glu	Tyr	Phe	Phe		
Leu 305	Glu	Tyr	Asn	Asn	lie 310	Gln	His	Leu	Phe	Ser 315	His	ser	Leu	His	Gly 320		
Leu	Phe	Asn	Val	Arg 325	Tyr	Leu	Asn	Leu	Lys 330	Arg	ser	Phe	Thr	Lys 335	Gln		
Ser	lie	Ser	Leu 340	Ala	Ser	Leu	Pro	Lys 345	lie	Asp	Asp	Phe	Ser 350	Phe	Gln		
Trp	Leu	Lys 355	cys	Leu	Glu	His	Leu 360	Asn	Met	Glu	Asp	Asn 365	Asp	lie	Pro		
Gly 370	lie	Lys	ser	Asn	Met	Phe 375	Thr	Gly	Leu	lie	Asn 380	Leu	Lys	Tyr	Leu		
Ser 385	Leu	ser	Asn	ser	Phe 390	Thr	ser	Leu	Arg	Thr 395	Leu	Thr	Asn	Glu	Thr 400		
Phe	Val	ser	Leu	Ala 405	His	Ser	Pro	Leu	His 410	lie	Leu	Asn	Leu	Thr 415	Lys		
Asn	Lys	lie	Ser 420	Lys	lie	Glu	Ser	Asp 425	Ala	Phe	Ser	Trp	Leu 430	Gly	His		
Leu	Glu	Val 435	Leu	Asp	Leu	Gly	Leu 440	Asn	Glu	lie	Gly	Gln 445	Glu	Leu	Thr		
Gly 450	Gln	Glu	Trp	Arg	Gly	Leu 455	Glu	Asn	lie	Phe	Glu 460	lie	Tyr	Leu	Ser		
Tyr 465	Asn	Lys	Tyr	Leu	Gln 470	Leu	Thr	Arg	Asn	Ser 475	Phe	Ala	Leu	Val	Pro 480		
Ser	Leu	Gln	Arg	Leu 485	Met	Leu	Arg	Arg	Val 490	Ala	Leu	Lys	Asn	Val 495	Asp		
Ser	Ser	Pro	ser 500	Pro	Phe	Gln	Pro	Leu 505	Arg	Asn	Leu	Thr	H e 510	Leu	Asp		
Leu	Ser	Asn 515	Asn	Asn	lie	Ala	Asn 520	lie	Asn	Asp	Asp	Met 525	Leu	Glu	Gly		
Leu	Glu 530	Lys	Leu	Glu	lie	Leu 535	Asp	Leu	Gln	His	Asn 540	Asn	Leu	Ala	Arg		

Leu 545	Trp	Lys	His	Ala	Asn 550	Pro	Gly	Gly	Pro	lie 555	Tyr	Phe	Leu	Lys	Gly 560
Leu	Ser	His	Leu	His 565	lie	Leu	Asn	Leu	Glu 570	Ser	Asn	Gly	Phe	Asp 575	Glu
lie	Pro	Val	Glu 580	Val	phe	Lys	Asp	Leu 585	Phe	Glu	Leu	Lys	lie 590	lie	Asp
Leu	Gly	Leu 595	Asn	Asn	Leu	Asn	Thr 600	Leu	Pro	Ala	Ser	val 605	Phe	Asn	Asn
Gin 610	val	Ser	Leu	Lys	Ser	Leu 615	Asn	Leu	Gin	Lys	Asn 620	Leu	lie	Thr	Ser
Val 625	Glu	Lys	Lys	val	Phe 630	Gly	Pro	Ala	Phe	Arg 635	Asn	Leu	Thr	Glu	Leu 640
Asp	Met	Arg	Phe	Asn 645	Pro	Phe	Asp	Cys	Thr 650	Cys	Glu	Ser	lie	Ala 655	Trp
Phe	Val	Asn	Trp 660	lie	Asn	Glu	Thr	His 665	Thr	Asn	lie	Pro	Glu 670	Leu	Ser
Ser	His	Tyr 675	Leu	Cys	Asn	Thr	Pro 680	Pro	His	Tyr	His	Gly 685	Phe	Pro	Val
Arg	Leu 690	Phe	Asp	Thr	Ser	Ser 695	cys	Lys	Asp	Ser	Ala 700	Pro	Phe	Glu	Leu
Phe 705	Phe	Met	lie	Asn	Thr 710	Ser	lie	Leu	Leu	lie 715	Phe	lie	Phe	lie	Val 720
Leu	Leu	lie	His	Phe 725	Glu	Gly	Trp	Arg	lie 730	ser	Phe	Tyr	Trp	Asn 735	val
ser	Val	His	Arg 740	Val	Leu	Gly	Phe	Lys 745	Glu	lie	Asp	Arg	Gin 750	Thr	Glu
Gin	Phe	Glu 755	Tyr	Ala	Ala	Tyr	lie 760	lie	His	Ala	Tyr	Lys 765	Asp	Lys	Asp
Trp 770	Val	Trp	Glu	His	Phe	Ser 775	Ser	Met	Glu	Lys	Glu 780	Asp	Gin	Ser	Leu
Lys 785	Phe	Cys	Leu	Glu	Glu 790	Arg	Asp	Phe	Glu	Ala 795	Gly	Val	Phe	Glu	Leu 800
Glu	Ala	lie	Val	Asn 805	Ser	lie	Lys	Arg	ser 810	Arg	Lys	lie	lie	Phe	Val 815

δ

lie Thr His His Leu Leu Lys Asp Pro Leu cys Lys Arg Phe Lys VaI
820 825 830

His His Ala VaI Gin Gin Ala lie Glu Gin Asn Leu Asp Ser lie lie
835 840 845

Leu VaI Phe Leu Glu Glu lie Pro Asp Tyr Lys Leu Asn His Ala Leu
850 855 860

Cys Leu Arg Arg Gly Met Phe Lys Ser His Cys lie Leu Asn Trp Pro
865 870 875 880

VaI Gin Lys Glu Arg lie Gly Ala Phe Arg His Lys Leu Gin VaI Ala
885 890 895

Leu Gly Ser Lys Asn ser VaI His
900

<210> 3
<211> 21
<212> DNA
<213> artificial sequence

<220>
<223> primer

<400> 3
ctcagaagat taccagccgc c 21

<210> 4
<211> 22
<212> DNA
<213> artificial sequence

<220>
<223> primer

<400> 4
ccattatgag acagatctaa tg 22

INTERNATIONAL SEARCH REPORT

International application No
PCT/IB2005/003641

A. CLASSIFICATION
C12N15/11 A61P35/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C12N A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal , WPI Data, PAJ, BIOSIS, EMBASE, PASCAL

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No
X	<p>ANDRE F (REPRINT) ET AL: "Targeting toll like receptor 3 by double stranded RNA in breast cancer: Results from in vitro studies and randomized trial."</p> <p>JOURNAL OF CLINICAL ONCOLOGY, (15 JUL 2004) VOL. 22, NO. 14, SUPP. 'S!, PP. 864S-864S. MA 9619. ISSN: 0732-183X. PB - AMER SOC CLINICAL ONCOLOGY, 330 JOHN CARLYLE ST, STE 300, ALEXANDRIA, VA 22314 USA., 15 July 2004 (2004-07-15), XP009051335</p> <p>the whole document</p> <p>----- -/--</p>	1-49

☒ Further documents are listed in the continuation of Box C

☒ See patent family annex

* Special categories of cited documents

"A" document defining the general state of the art which is not considered to be of particular relevance
 "E" earlier document but published on or after the international filing date
 "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
 "O" document referring to an oral disclosure, use, exhibition or other means
 "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
 "X¹" document of particular relevance, the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
 "Y" document of particular relevance, the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
 "&" document member of the same patent family

Date of the actual completion of the international search

9 February 2006

Date of mailing of the international search report

27/02/2006

Name and mailing address of the ISA/

European Patent Office, P B 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel (+31-70) 340-2040, Tx 31 651 epo nl,
 Fax (+31-70) 340-3016

Authorized officer

Fayos , C

INTERNATIONAL SEARCH REPORT

International application No
PCT/IB2005/003641

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No
X	<p>WHITMORE MARK M ET AL: "Synergi stic activation of innate immunity by double-stranded RNA and CpG DNA promotes enhanced antitumor activity"</p> <p>CANCER RESEARCH, vol . 64, no. 16, 15 August 2004 (2004-08-15) , pages 5850-5860, XP009051341 ISSN: 0008-5472 page 5850, col umn 2, last paragraph; figure 5 page 5853, col umn 1, paragraphs 2,3 page 5855, col umn 1, last paragraph; figure 5 page 5857, col umn 2 page 5858, col umn 2, paragraphs 2,3</p>	1-49
X	<p>EP 0 281 380 A (HEM RESEARCH, INC ; HEM PHARMACEUTICALS CORP) 7 September 1988 (1988-09-07) cited in the appl icati on claim 1 page 3, line 1 - line 45 figures 5,6</p>	1-49
X	<p>EP 0 113 162 A (CARTER, WILLIAM ALVIN; TS'O, PAUL O.P) 11 July 1984 (1984-07-11) cited in the appl icati on claims 1-14 page 1, line 19 - line 28 page 9, line 17 - page 11, line 32</p>	1-49
Y	<p>ALEXOPOULOU L ET AL: "Recogniti on of double-stranded RNA and acti vati on of NF-kB by tol l-like receptor 3"</p> <p>NATURE, MACMILLAN JOURNALS LTD. LONDON , GB, vol . 413, 18 October 2001 (2001-10-18) , pages 732-738, XP002968529 ISSN: 0028-0836 the whole document</p>	1-49
X	<p>us 2003/166001 AI (LIPFORD GRAYSON B) 4 September 2003 (2003-09-04)</p>	21-30
Y	<p>the whole document</p>	1-49

-/--

INTERNATIONAL SEARCH REPORT

International application No
PCT/IB2005/003641

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No
P,X	<p>M ET AL: "The rationale for combined chemo/immunotherapy using a Toll-I like receptor 3 (TLR3) agonist and tumour-derived exosomes in advanced ovarian cancer"</p> <p>VACCINE, BUTTERWORTH SCIENTIFIC. GUILDFORD, GB, vol. 23, no. 17-18, 18 March 2005 (2005-03-18), pages 2374-2378, XP004777559</p> <p>ISSN: 0264-410X</p> <p>the whole document</p> <p>-----</p>	1-49

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/IB2005/003641

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0281380	A	07-09-1988	AT 130760 T	15-12-1995
			AU 1256588 A	01-09-1988
			AU 1618992 A	06-08-1992
			AU 6348398 A	11-06-1998
			CA 1336810 C	29-08-1995
			CN 1032296 A	12-04-1989
			DE 3854726 D1	11-01-1996
			DE 3854726 T2	25-07-1996
			DK 113188 A	04-09-1988
			ES 2082749 T3	01-04-1996
			FI 880961 A	04-09-1988
			IE 75895 B1	24-09-1997
			IL 85603 A	13-05-1993
			JP 1980506 C	17-10-1995
			JP 7013024 B	15-02-1995
			JP 63295514 A	01-12-1988
			KR 9608009 B1	19-06-1996
			NO 880946 A	05-09-1988
			NZ 223720 A	26-07-1991
			PH 24467 A	18-07-1990
			PT 86879 A	01-04-1988
			SU 1836103 A3	23-08-1993
			ZA 8801512 A	30-11-1988
EP 0113162	A	11-07-1984	DE 3380200 D1	24-08-1989
			JP 1768960 C	30-06-1993
			JP 4054649 B	31-08-1992
			JP 59134735 A	02-08-1984
US 2003166001	A1	04-09-2003	NONE	